

Riesz bases associated with regular representations of semidirect product groups

Antonio G. García* and Gerardo Pérez-Villalón†

* Departamento de Matemáticas, Universidad Carlos III de Madrid, Avda. de la Universidad 30, 28911 Leganés-Madrid, Spain.

† Departamento de Matemática Aplicada a las Tecnologías de la Información y las Comunicaciones, E.T.S.I.T., Universidad Politécnica de Madrid, Avda. Complutense 30, 28040 Madrid, Spain.

Abstract

This work is devoted to the study of Bessel and Riesz systems of the type $\{L_\gamma f\}_{\gamma \in \Gamma}$ obtained from the action of the left regular representation L_γ of a discrete non abelian group Γ which is a semidirect product, on a function $f \in \ell^2(\Gamma)$. The main features about these systems can be conveniently studied by means of a simple matrix-valued function $\mathbf{F}(\xi)$. These systems allow to derive sampling results in principal Γ -invariant spaces, i.e., spaces obtained from the action of the group Γ on a element of a Hilbert space. Since the systems $\{L_\gamma f\}_{\gamma \in \Gamma}$ are closely related to convolution operators, a connection with C^* -algebras is also established.

Keywords: Semidirect product of groups; left regular representation of a group; dual Riesz bases; sampling expansions.

AMS: 42C15; 20H15; 94A20.

1 Introduction

This work is devoted to the study of a characterization as Riesz bases, together with some sampling applications, of systems $\{L_\gamma f\}_{\gamma \in \Gamma}$ obtained from the left regular representation of a discrete non abelian group Γ , that is, $L_\gamma f(\eta) := f(\gamma^{-1}\eta)$, $\eta, \gamma \in \Gamma$, where f denotes a fixed element in the Hilbert space $\ell^2(\Gamma)$. Throughout the paper the group $\Gamma := N \rtimes_\sigma H$ is the semidirect product of two groups: a discrete abelian group N and a finite group H ; the subscript σ denotes the action of the group H on the group N . Some important examples of non abelian groups such as dihedral groups, infinite dihedral group or crystallographic groups are semidirect products with these characteristics.

In addition to the intrinsic importance of the left regular representation $\gamma \mapsto L_\gamma \in \mathcal{U}(\ell^2(\Gamma))$ in representation theory of groups, the systems $\{L_\gamma f\}_{\gamma \in \Gamma}$ arising from the left regular representation of Γ are relevant in applications; for instance they appear in sampling theory. In

*E-mail: agarcia@math.uc3m.es

†E-mail: gperez@euitt.upm.es

fact, in the present paper we deal with two types of samples where these systems have an important role:

Firstly, given a unitary representation $\Gamma \ni \gamma \mapsto U(\gamma) \in \mathcal{U}(\mathcal{H})$ of the group Γ on a separable Hilbert space \mathcal{H} , for a fixed $\varphi \in \mathcal{H}$ we consider the subspace of \mathcal{H}

$$\mathcal{A}_\varphi = \left\{ \sum_{\gamma \in \Gamma} \mathbf{a}(\gamma) U(\gamma) \varphi : \mathbf{a} = \{\mathbf{a}(\gamma)\}_{\gamma \in \Gamma} \in \ell^2(\Gamma) \right\}$$

For a fixed $\psi \in \mathcal{H}$, which does not necessarily belong to \mathcal{A}_φ , we can define for each $f \in \mathcal{A}_\varphi$ its samples

$$\mathcal{L}_\psi f(\gamma) := \langle f, U(\gamma) \psi \rangle_{\mathcal{H}}, \quad \gamma \in \Gamma.$$

These samples give average sampling in classical shift-invariant subspaces of $L^2(\mathbb{R})$ (see, for instance, Refs. [1, 9, 11, 17]). As we will see in Section 4, there exists $\mathbf{f}_\psi \in \ell^2(\Gamma)$ such that, for each $f \in \mathcal{A}_\varphi$, we get $\mathcal{L}_\psi f(\gamma) = \langle \mathbf{a}, L_\gamma \mathbf{f}_\psi \rangle_{\ell^2(\Gamma)}$, $\gamma \in \Gamma$, being $\mathbf{a} \in \ell^2(\Gamma)$ the coefficients sequence of $f \in \mathcal{A}_\varphi$.

Secondly, when $\mathcal{H} = L^2(\mathbb{R}^d)$, for a fixed point $p \in \mathbb{R}^d$ we consider, for any $f \in \mathcal{A}_\varphi$ the samples

$$\mathcal{L}_p f(\gamma) := [U(\gamma^{-1})f](p), \quad \gamma \in \Gamma.$$

Again, there exists $\mathbf{f}_p \in \ell^2(\Gamma)$ such that, for each $f \in \mathcal{A}_\varphi$, we obtain the expression for these samples $\mathcal{L}_p f(\gamma) = \langle \mathbf{a}, L_\gamma \mathbf{f}_p \rangle_{\ell^2(\Gamma)}$, $\gamma \in \Gamma$. This situation englobes the case when we are dealing with pointwise samples in classical shift-invariant subspaces of $L^2(\mathbb{R})$ [11, 16, 17, 23, 27].

Needless to say that a feasible characterization of the system $\{L_\gamma \mathbf{f}\}_{\gamma \in \Gamma}$ as a Riesz basis for $\ell^2(\Gamma)$ along with the search of its dual Riesz basis will play a crucial role in obtaining an interpolatory sampling formula allowing the recovery of any $f \in \mathcal{A}_\varphi$ from the given data sampling $\{\mathcal{L}_\psi f(\gamma)\}_{\gamma \in \Gamma}$ or $\{\mathcal{L}_p f(\gamma)\}_{\gamma \in \Gamma}$. Some Γ -invariant spaces \mathcal{A}_φ of special relevance are those appearing in composite wavelet theory. These wavelets allow many more locations, scales and directions than the classical ones. They have been studied in the last few years; see, for instance, Refs. [18, 19, 20, 21].

In order to get a suitable characterization of when the system $\{L_\gamma \mathbf{f}\}_{\gamma \in \Gamma}$ is a Riesz basis for $\ell^2(\Gamma)$, we briefly describe the mathematical techniques used in this paper. Since, for $\mathbf{a}, \mathbf{f} \in \ell^2(\Gamma)$, we have

$$\sum_{\eta \in \Gamma} \mathbf{a}(\eta) L_\eta \mathbf{f} = \mathbf{a} * \mathbf{f} \quad \text{and} \quad \langle \mathbf{a}, L_\gamma \mathbf{f} \rangle_{\ell^2(\Gamma)} = (\mathbf{a} * \mathbf{f}^*)(\gamma),$$

where \mathbf{f}^* denotes the involution in $\ell^2(\Gamma)$ of \mathbf{f} , we can use techniques of linear time-invariant (LTI) systems. In fact, the given characterization will be described in terms of a matrix-valued function $\mathbf{F}(\xi)$ which turns out to be the transfer matrix of a multi-input multi-output (MIMO) system. Although, our group $\Gamma = N \rtimes_\sigma H$ is not abelian and a classical Fourier analysis is not directly applicable, the MIMO system formalism will allow us to make use of the Fourier transform on the locally compact abelian (LCA) group N ; in fact, $\mathbf{F}(\xi)$ is defined for ξ in \widehat{N} , the dual group of characters ξ of N .

As the above equalities show, the study of the systems $\{L_\gamma \mathbf{f}\}_{\gamma \in \Gamma}$ is related to convolution algebras on Γ , and specially, to the representation of the group C^* -algebra of Γ , $C^*(\Gamma)$, given

in Ref. [25]. In the last section we show this relationship and, in so doing, we provide a convolution C^* -algebra larger than $C^*(\Gamma)$ and suitable for the present context.

The paper is organized as follows: Section 2 provides the mathematical setting needed throughout the paper giving the keys for different approaches, together with some lemmata used in the sequel. Section 3 includes the main theoretical results in the paper: A characterization of when $\{L_\gamma f\}_{\gamma \in \Gamma}$ is a Bessel sequence or a Riesz basis for $\ell^2(\Gamma)$ is respectively proved in Theorems 5 and 6; in particular the orthonormal basis case is considered (Corollary 7). The dual Riesz basis of $\{L_\gamma f\}_{\gamma \in \Gamma}$, which has the same form $\{L_\gamma g\}_{\gamma \in \Gamma}$, is also obtained (Corollary 8). Closing the section, an example of Riesz bases associated to the infinite dihedral group D_∞ is proposed. Section 4 is devoted to a sampling application of the results in Section 3; thus, an abstract sampling result is obtained (Theorem 9) in a principal U -invariant subspace of a Hilbert space \mathcal{H} . An example using crystallographic groups illustrates the sampling results, where we consider pointwise samples as well as average samples. Finally, in Section 5 a C^* -algebras connection is also exhibited.

2 The mathematical setting

The aim of this section is twofold: Firstly, to provide a brief on the needed mathematical preliminaries and, secondly, to establish the mathematical setting which will give us the keys and tools to get the main aim in the paper, i.e., a suitable characterization of the system $\{L_\gamma f\}_{\gamma \in \Gamma}$ as a Riesz basis for $\ell^2(\Gamma)$.

We begin stating the main facts concerning the group Γ as a semidirect product of groups. Let $(N, +)$ be a discrete abelian group, (H, \cdot) a finite group of order κ , and a homomorphism $\sigma : H \mapsto \text{Aut}(N)$ referred as the action of the group H on the group N . Its *semidirect product* $\Gamma := N \rtimes_\sigma H$ is the group whose elements are the pairs $(n, h) \in N \times H$ with multiplication rule

$$(n, h)(m, l) := (n + \sigma(h)(m), hl), \quad n, m \in N \text{ and } h, l \in H.$$

In particular, the identity element in Γ is $e_\Gamma = (0_N, 1_H)$ and $(n, h)^{-1} = (-\sigma(h^{-1})(n), h^{-1})$, $(n, h) \in \Gamma$. Notice that, unless $\sigma(h)$ equals the identity for $h \in H$, the group Γ is not abelian.

An important example of semidirect product of groups that will be used in this paper is the *crystallographic group* $\Gamma_{\mathcal{M}, H} = \mathcal{M}\mathbb{Z}^d \rtimes_\sigma H$, where \mathcal{M} is a non-singular $d \times d$ matrix and H is a finite subgroup of $O(d)$, the orthogonal group of order d , such that $A(\mathcal{M}\mathbb{Z}^d) = \mathcal{M}\mathbb{Z}^d$ for all $A \in H$. Here $\sigma(A)x = Ax$ for $A \in H$ and $x \in \mathbb{R}^d$. The *infinite dihedral group* $D_\infty := \mathbb{Z} \rtimes_\sigma \mathbb{Z}_2$, where $\sigma(1)(n) = n$ and $\sigma(-1)(n) = -n$ for each $n \in \mathbb{Z}$, is a unidimensional crystallographic group.

Throughout the paper we denote by greek letters γ, η, \dots or as $(n, h), (m, l), \dots$ the elements in Γ . The *left regular representation* of the group Γ on $\ell^2(\Gamma)$ is given by

$$L_\gamma f(\eta) := f(\gamma^{-1}\eta), \quad \eta, \gamma \in \Gamma \text{ and } f \in \ell^2(\Gamma).$$

Note that, for each $f \in \ell^2(\Gamma)$, the *synthesis operator* Λ_f of the system $\{L_\gamma f\}_{\gamma \in \Gamma}$ is a convolution operator. Indeed, for any $\mathbf{a} \in \ell^2(\Gamma)$

$$\Lambda_f \mathbf{a}(\gamma) := \sum_{\eta \in \Gamma} \mathbf{a}(\eta) L_\eta f(\gamma) = \sum_{\eta \in \Gamma} \mathbf{a}(\eta) f(\eta^{-1}\gamma) = (\mathbf{a} * f)(\gamma), \quad \gamma \in \Gamma.$$

Recall that the above definition gives a bounded linear operator $\Lambda_f : \ell^2(\Gamma) \mapsto \ell^2(\Gamma)$ if and only if the system $\{L_\gamma f\}_{\gamma \in \Gamma}$ is a Bessel sequence [8, Theorem 3.2.3] (see also Theorem 5 infra). Our study is based on the following representation of $\Lambda_f \mathbf{a}$:

Lemma 1. *Given $f, \mathbf{a} \in \ell^2(\Gamma)$, $\Lambda_f \mathbf{a}$ is represented as:*

$$\Lambda_f \mathbf{a}(n, h) = \sum_{l \in H} (\mathbf{a}_l *_{N} f_{h,l})(n), \quad (n, h) \in \Gamma, \quad (1)$$

where $\mathbf{a}_l(n) := \mathbf{a}(n, l)$, $f_{h,l}(n) := f[(0, l)^{-1}(n, h)]$, and $*_{N}$ denotes the convolution on the abelian group N , i.e., $(a *_{N} b)(n) := \sum_{m \in N} a(m) b(n - m)$.

Proof. Having in mind that $(m, l)^{-1} = (0, l)^{-1}(-m, 1)$, we get

$$\Lambda_f \mathbf{a}(n, h) = \sum_{l \in H} \sum_{m \in N} \mathbf{a}(m, l) f[(m, l)^{-1}(n, h)] = \sum_{l \in H} \sum_{m \in N} \mathbf{a}(m, l) f[(0, l)^{-1}(n - m, h)],$$

and representation (1) holds. \square

According to expression (1), the operator Λ_f can be seen as a linear time-invariant system:

$$[\mathbf{a}_l]_{l \in H} \in \ell_{\kappa}^2(N) \mapsto \left[\sum_{l \in H} \mathbf{a}_l *_{N} f_{h,l} \right]_{h \in H} \quad (2)$$

where $\ell_{\kappa}^2(N) := \ell^2(N) \times \dots \times \ell^2(N)$ (κ times). In signal processing jargon this type of system is called a multi-input, multi-output (MIMO) linear time-invariant system; it can be effectively analyzed by using the Fourier transform.

Since N is a discrete abelian group, we can use the Fourier transform on N defined by

$$\widehat{\mathbf{a}}(\xi) = \sum_{n \in N} \mathbf{a}(n) \langle -n, \xi \rangle, \quad \xi \in \widehat{N},$$

for any $\mathbf{a} \in \ell^1(N)$, and extended to $\ell^2(N)$ as a unitary operator between $\ell^2(N)$ and $L^2(\widehat{N})$ where \widehat{N} denotes the dual group of characters (see, for instance, Ref.[10] for the details).

Next lemma will be needed in taking the Fourier transform in Eq. (1). It also gives a condition so that the output in (2) belongs to $\ell_{\kappa}^2(N)$.

Lemma 2. *Let $\mathbf{a}, \mathbf{b} \in \ell^2(N)$ such that the product $\widehat{\mathbf{a}}(\xi) \widehat{\mathbf{b}}(\xi) \in L^2(\widehat{N})$. Then the convolution $\mathbf{a} *_{N} \mathbf{b} \in \ell^2(N)$ and*

$$(\widehat{\mathbf{a} *_{N} \mathbf{b}})(\xi) = \widehat{\mathbf{a}}(\xi) \widehat{\mathbf{b}}(\xi), \quad a.e. \xi \in \widehat{N}.$$

Proof. By using Plancherel theorem [10, Theorem 4.25] and denoting $\widetilde{\mathbf{b}}(n) = \overline{\mathbf{b}(-n)}$, we obtain

$$\begin{aligned} (\mathbf{a} *_{N} \mathbf{b})(n) &= \sum_{m \in N} \mathbf{a}(m) \mathbf{b}(n - m) = \langle \mathbf{a}, \widetilde{\mathbf{b}}(\cdot - n) \rangle_{\ell^2(N)} = \langle \widehat{\mathbf{a}}, \widehat{\widetilde{\mathbf{b}}(\cdot - n)} \rangle_{L^2(\widehat{N})} \\ &= \int_{\widehat{N}} \widehat{\mathbf{a}}(\xi) \overline{\widehat{\widetilde{\mathbf{b}}(\cdot - n)}(\xi)} d\mu_{\widehat{N}}(\xi) = \int_{\widehat{N}} \widehat{\mathbf{a}}(\xi) \widehat{\mathbf{b}}(\xi) \overline{\langle -n, \xi \rangle} d\mu_{\widehat{N}}(\xi). \end{aligned}$$

Since $\{\langle -n, \xi \rangle\}_{n \in N}$ is an orthonormal basis for $L^2(\widehat{N})$ [10, Theorem 4.26] and we have assumed that $\widehat{\mathbf{a}}(\xi) \widehat{\mathbf{b}}(\xi) \in L^2(\widehat{N})$, we obtain that $\mathbf{a} *_N \mathbf{b} \in \ell^2(N)$ and

$$\widehat{\mathbf{a}}(\xi) \widehat{\mathbf{b}}(\xi) = \sum_{n \in N} (\mathbf{a} *_N \mathbf{b})(n) \langle -n, \xi \rangle, \quad \text{a.e. } \xi \in \widehat{N}.$$

□

By taking the N -Fourier transform in the second term of expression (1) we obtain the so called *transfer matrix* of the MIMO system (2). This motivates the following definition:

Definition 1. For each $\mathbf{f} \in \ell^2(\Gamma)$ we introduce its associated transfer matrix as the $\kappa \times \kappa$ matrix-valued function \mathbf{F} defined on \widehat{N} as

$$\mathbf{F}(\xi) = [\widehat{\mathbf{f}}_{h,l}(\xi)]_{h,l \in H} \quad \text{where } \mathbf{f}_{h,l}(n) = \mathbf{f}[(0,l)^{-1}(n,h)], \quad n \in N. \quad (3)$$

The *involution* in $\ell^2(\Gamma)$ and in $\ell^2(N)$ are denoted, respectively, by

$$\mathbf{f}^*(\gamma) = \overline{\mathbf{f}(\gamma^{-1})}, \quad \gamma \in \Gamma \quad \text{and} \quad \widetilde{\mathbf{f}}_{h,l}(n) = \overline{\mathbf{f}_{h,l}(-n)}, \quad n \in N.$$

The role of the conjugate transpose matrix-valued function $\mathbf{F}^*(\xi)$ is also well understood realizing that it is the transfer matrix of the system

$$[\mathbf{a}_l]_{l \in H} \mapsto \left[\sum_{l \in H} \mathbf{a}_l *_N \widetilde{\mathbf{f}}_{l,h} \right]_{h \in H}$$

which represents the *analysis operator* of the sequence $\{L_\gamma \mathbf{f}\}_{\gamma \in \Gamma}$. Namely, for \mathbf{f}, \mathbf{a} in $\ell^2(\Gamma)$, we have

$$\mathcal{A}_\mathbf{f}(n, h) := \langle \mathbf{a}, L_{(n,h)} \mathbf{f} \rangle_{\ell^2(\Gamma)} = (\mathbf{a} * \mathbf{f}^*)(n, h) = \sum_{l \in H} (\mathbf{a}_l *_N \widetilde{\mathbf{f}}_{l,h})(n). \quad (4)$$

Indeed, equality

$$\sum_{l \in H} \sum_{m \in N} \mathbf{a}(m, l) \overline{\mathbf{f}[(n, h)^{-1}(m, l)]} = \sum_{l \in H} \sum_{m \in N} \mathbf{a}(m, l) \overline{\mathbf{f}[(0, h)^{-1}(m - n, l)]}$$

yields (4). Recall that the analysis operator is the adjoint operator of the synthesis operator [8, Lemma 3.2.1]; in other words, whenever the system $\{L_\gamma \mathbf{f}\}_{\gamma \in \Gamma}$ is a Bessel sequence for $\ell^2(\Gamma)$, operator $\mathcal{A}_\mathbf{f}$ in (4) is the adjoint operator of $\Lambda_\mathbf{f}$, i.e., $\mathcal{A}_\mathbf{f} = \Lambda_\mathbf{f}^*$.

In our context, we will need the transform \mathcal{T}_Γ given in the following lemma, where $L_\kappa^2(\widehat{N})$ denotes the product Hilbert space $L^2(\widehat{N}) \times \dots \times L^2(\widehat{N})$ (κ times):

Lemma 3. The linear map $\mathcal{T}_\Gamma : \ell^2(\Gamma) \rightarrow L_\kappa^2(\widehat{N})$ defined by $\mathcal{T}_\Gamma \mathbf{a} := [\widehat{\mathbf{a}}_h]_{h \in H}$, where $\mathbf{a}_h(n) = \mathbf{a}(n, h)$, $(n, h) \in \Gamma$, is a unitary operator.

Proof. The map \mathcal{T}_Γ is surjective since the N -Fourier transform is a unitary operator between $\ell^2(N)$ and $L^2(\widehat{N})$. It is also an isometry since, for each $\mathbf{a}, \mathbf{b} \in \ell^2(\Gamma)$, we have

$$\langle \mathbf{a}, \mathbf{b} \rangle_{\ell^2(\Gamma)} = \sum_{h \in H} \langle \mathbf{a}_h, \mathbf{b}_h \rangle_{\ell^2(N)} = \sum_{h \in H} \langle \widehat{\mathbf{a}}_h, \widehat{\mathbf{b}}_h \rangle_{L^2(\widehat{N})} = \langle \mathcal{T}_\Gamma \mathbf{a}, \mathcal{T}_\Gamma \mathbf{b} \rangle_{L_\kappa^2(\widehat{N})}.$$

□

The above lemma, related to the abstract version of the Zak transform (see, for instance, Refs.[3, 12]), says that any $\mathbf{a} \in \ell^2(\Gamma)$ is completely determined by the Fourier transform of its κ phases \mathbf{a}_h , $h \in H$.

With a view to built the matrix-valued function $\mathbf{F}(\xi)$ and the vector-valued function $\mathcal{T}_\Gamma \mathbf{a}(\xi)$, indexed by the elements of H , we order the κ elements of H such that the first element is 1_H , the identity element of H . Thus the first column of $\mathbf{F}(\xi)$ is $\mathcal{T}_\Gamma \mathbf{f}(\xi)$. Notice also that its l -column is $\mathcal{T}_\Gamma L_{(0_N, l)} \mathbf{f}(\xi)$. Hence, for $\xi \in \widehat{N}$ matrix $\mathbf{F}(\xi)$ is a redundant matrix, similar to what happens with the modulation matrix in wavelet or in filter bank theory.

In next result, we obtain a representation for the analysis and synthesis operators, Eqs. (1) and (4) respectively, in the \mathcal{T}_Γ domain:

Theorem 4. *Assume that $\mathbf{f}, \mathbf{a} \in \ell^2(N)$ and that the products $\widehat{\mathbf{a}}_l(\xi) \widehat{\mathbf{f}}_{h, h'}(\xi) \in L^2(\widehat{N})$ for all $l, h, h' \in H$. Then*

$$\mathcal{T}_\Gamma \Lambda_{\mathbf{f}} \mathbf{a}(\xi) = \mathbf{F}(\xi) \mathcal{T}_\Gamma \mathbf{a}(\xi) \quad \text{and} \quad \mathcal{T}_\Gamma \mathcal{A}_{\mathbf{f}} \mathbf{a}(\xi) = \mathbf{F}^*(\xi) \mathcal{T}_\Gamma \mathbf{a}(\xi), \quad \text{a.e. } \xi \in \widehat{N}.$$

Besides, on the assumption that $\mathbf{b} = \mathbf{a} * \mathbf{f}$ then

$$\mathbf{B}(\xi) = \mathbf{F}(\xi) \mathbf{A}(\xi), \quad \text{a.e. } \xi \in \widehat{N}, \quad (5)$$

where $\mathbf{A}(\xi)$ and $\mathbf{B}(\xi)$ are the transfer matrices associated to \mathbf{a} and \mathbf{b} defined in (3).

Proof. By taking the N -Fourier transform in equalities (1) and (4), and having in mind Lemma 2 we obtain that $\mathcal{T}_\Gamma \Lambda_{\mathbf{f}} \mathbf{a}(\xi) = \mathbf{F}(\xi) \mathcal{T}_\Gamma \mathbf{a}(\xi)$ and $\mathcal{T}_\Gamma \mathcal{A}_{\mathbf{f}} \mathbf{a}(\xi) = \mathbf{F}^*(\xi) \mathcal{T}_\Gamma \mathbf{a}(\xi)$. Concerning the second part, the l -column of $\mathbf{A}(\xi)$ is $\mathcal{T}_\Gamma(L_{(0_N, l)} \mathbf{a})$ and the l -column of $\mathbf{B}(\xi)$ is

$$\mathcal{T}_\Gamma(L_{(0_N, l)} \mathbf{b})(\xi) = \mathcal{T}_\Gamma(L_{(0_N, l)}[\mathbf{a} * \mathbf{f}])(\xi) = \mathcal{T}_\Gamma([L_{(0_N, l)} \mathbf{a}] * \mathbf{f})(\xi) = \mathbf{F}(\xi) \mathcal{T}_\Gamma(L_{(0_N, l)} \mathbf{a})(\xi),$$

that is, the l -column of $\mathbf{F}(\xi) \mathbf{A}(\xi)$. □

A similar formula to (5) is obtained in Ref. [25] for functions in the C^* -algebra of the group Γ , even for a non discrete group N ; see Section 5 below. It can be also found in Ref. [21] for functions in $\ell^1(\Gamma)$, being Γ a crystallographic group.

3 Riesz bases for $\ell^2(\Gamma)$ generated by the left regular representation of Γ

In what follows we will assume that the non abelian group Γ is the semidirect product $N \rtimes_\sigma H$, where $(N, +)$ is a discrete abelian group and (H, \cdot) is a finite group of order κ . For a fixed $\mathbf{f} \in \ell^2(\Gamma)$, this section is devoted to give a characterization of the sequence $\{L_\gamma \mathbf{f}\}_{\gamma \in \Gamma}$ as a Riesz basis for $\ell^2(\Gamma)$ in terms of the associated matrix-valued function $\mathbf{F}(\xi)$ introduced in (3).

Theorem 5. *For $\mathbf{f} \in \ell^2(\Gamma)$, let $\mathbf{F}(\xi)$ be its associated transfer matrix defined in (3). Then, the system $\{L_\gamma \mathbf{f}\}_{\gamma \in \Gamma}$ is a Bessel sequence for $\ell^2(\Gamma)$ if and only if the entries of the matrix-valued function $\mathbf{F}(\xi)$ belong to $L^\infty(\widehat{N})$. In this case the optimal Bessel bound is given by*

$$B_{\mathbf{f}} := \operatorname{ess\,sup}_{\xi \in \widehat{N}} \lambda_{\max}[\mathbf{F}^*(\xi) \mathbf{F}(\xi)],$$

where λ_{\max} denotes the largest eigenvalue of $\mathbf{F}^*(\xi) \mathbf{F}(\xi)$.

Proof. Having in mind the equivalence between the spectral and the Frobenius norms for matrices [14], we deduce that $B_f < \infty$ if and only if the entries $\widehat{f}_{h,l}$ of $\mathbf{F}(\xi)$ belong to $L^\infty(\widehat{N})$.

Whenever $\mathbf{a} \in \ell^2(N)$ with $\widehat{a}_l(\xi) \widehat{f}_{h,h'}(\xi) \in L^2(\widehat{N})$, $l, h, h' \in H$, by using Lemma 3 and Theorem 4 we obtain that

$$\left\| \sum_{\gamma \in \Gamma} \mathbf{a}(\gamma) L_\gamma \mathbf{f} \right\|_{\ell^2(\Gamma)}^2 = \|\Lambda_f \mathbf{a}\|_{\ell^2(\Gamma)}^2 = \|\mathcal{T}_\Gamma \Lambda_f \mathbf{a}\|_{L_k^2(\widehat{N})}^2 = \left\| \mathbf{F}(\cdot) \mathcal{T}_\Gamma \mathbf{a}(\cdot) \right\|_{L_k^2(\widehat{N})}^2,$$

and then,

$$\left\| \sum_{\gamma \in \Gamma} \mathbf{a}(\gamma) L_\gamma \mathbf{f} \right\|_{\ell^2(\Gamma)}^2 = \int_{\widehat{N}} [\mathcal{T}_\Gamma \mathbf{a}(\xi)]^* \mathbf{F}^*(\xi) \mathbf{F}(\xi) \mathcal{T}_\Gamma \mathbf{a}(\xi) d\mu_{\widehat{N}}(\xi). \quad (6)$$

Hence, using Lemma 3, we obtain that, whenever $\mathbf{a} \in \ell^2(N)$ and $\widehat{f}_{h,l}(\xi) \in L^\infty(\widehat{N})$, $l, h \in H$, we have

$$\begin{aligned} \left\| \sum_{\gamma \in \Gamma} \mathbf{a}(\gamma) L_\gamma \mathbf{f}(\gamma) \right\|_{\ell^2(\Gamma)}^2 &\leq \int_{\widehat{N}} \lambda_{\max}[\mathbf{F}^*(\xi) \mathbf{F}(\xi)] \|\mathcal{T}_\Gamma \mathbf{a}(\xi)\|^2 d\xi \\ &\leq B_f \int_{\widehat{N}} \|\mathcal{T}_\Gamma \mathbf{a}(\xi)\|^2 d\xi = B_f \|\mathcal{T}_\Gamma \mathbf{a}\|_{L_k^2(\widehat{N})}^2 = B_f \|\mathbf{a}\|_{\ell^2(\Gamma)}^2. \end{aligned}$$

Consequently, if $\widehat{f}_{h,l}(\xi) \in L^\infty(\widehat{N})$, $h, l \in H$, or equivalently $B_f < \infty$, then $\{L_\gamma \mathbf{f}\}_{\gamma \in \Gamma}$ is a Bessel sequence with bound B_f .

For any number $J < B_f$, there exists a subset $\Omega \subset \widehat{N}$ with positive measure such that $\lambda_{\max}[\mathbf{F}^*(\xi) \mathbf{F}(\xi)] > J$ for $\xi \in \Omega$. Let $\mathbf{a} \in \ell^2(\Gamma)$ such that $\mathcal{T}_\Gamma \mathbf{a}(\xi)$ is equal to 0 when $\xi \notin \Omega$, and it is equal to a unitary eigenvector of $\mathbf{F}^*(\xi) \mathbf{F}(\xi)$ corresponding to the eigenvalue $\lambda_{\max}[\mathbf{F}^*(\xi) \mathbf{F}(\xi)]$ when $\xi \in \Omega$. By using (6) we obtain

$$\left\| \sum_{\gamma \in \Gamma} \mathbf{a}(\gamma) L_\gamma \mathbf{f}(\gamma) \right\|_{\ell^2(\Gamma)}^2 = \int_{\widehat{N}} \lambda_{\max}[\mathbf{F}^*(\xi) \mathbf{F}(\xi)] \|\mathcal{T}_\Gamma \mathbf{a}(\xi)\|^2 d\xi \geq J \int_{\widehat{N}} \|\mathcal{T}_\Gamma \mathbf{a}(\xi)\|^2 d\xi = J \|\mathbf{a}\|_{\ell^2(\Gamma)}^2.$$

Therefore, if $\{L_\gamma \mathbf{f}\}_{\gamma \in \Gamma}$ is Bessel sequence then $B_f < \infty$, or equivalently $\widehat{f}_{h,l}(\xi) \in L^\infty(\widehat{N})$, $h, l \in H$. Moreover, the constant B_f is the optimal Bessel bound. \square

Theorem 6. Consider $\mathbf{f} \in \ell^2(\Gamma)$ and its associated transfer matrix $\mathbf{F}(\xi)$ given in (3). The following statements are equivalent:

- (a) The system $\{L_\gamma \mathbf{f}\}_{\gamma \in \Gamma}$ is a Riesz basis for $\ell^2(\Gamma)$.
- (b) The entries of the matrix-valued function $\mathbf{F}(\xi)$ belongs to $L^\infty(\widehat{N})$, and $\text{ess inf}_{\xi \in \widehat{N}} |\det \mathbf{F}(\xi)| > 0$.

In this case the optimal Riesz bounds are given by

$$A_f := \text{ess inf}_{\xi \in \widehat{N}} \lambda_{\min}[\mathbf{F}^*(\xi) \mathbf{F}(\xi)] \quad \text{and} \quad B_f := \text{ess sup}_{\xi \in \widehat{N}} \lambda_{\max}[\mathbf{F}^*(\xi) \mathbf{F}(\xi)].$$

Moreover, its dual Riesz basis is $\{L_\gamma \mathbf{g}\}_{\gamma \in \Gamma}$ where \mathbf{g} is the unique element in $\ell^2(\Gamma)$ satisfying

$$\mathbf{F}^*(\xi) \mathcal{T}_\Gamma \mathbf{g}(\xi) = [1 \quad 0 \quad \dots \quad 0]^\top, \quad \text{a.e. } \xi \in \widehat{N}, \quad (7)$$

where \mathcal{T}_Γ is defined in Lemma 3. Equivalently, $\mathbf{G}(\xi) = (\mathbf{F}^*(\xi))^{-1}$, a.e. $\xi \in \widehat{N}$ where $\mathbf{G}(\xi)$ is the transfer matrix associated to \mathbf{g} .

Proof. First of all, note that for any $\kappa \times \kappa$ hermitian matrix \mathbf{M} we have that

$$\lambda_{\min}^{\kappa}(\mathbf{M}) \leq \det \mathbf{M} = \lambda_{\min}(\mathbf{M}) \cdots \lambda_{\max}(\mathbf{M}) \leq \lambda_{\min}(\mathbf{M}) \lambda_{\max}^{\kappa-1}(\mathbf{M}).$$

Using these inequalities for $\mathbf{M} = \mathbf{F}^*(\xi)\mathbf{F}(\xi)$ we obtain that

$$A_{\mathbf{f}}^{\kappa} \leq \operatorname{ess\,inf}_{\xi \in \widehat{N}} |\det \mathbf{F}(\xi)|^2 \leq A_{\mathbf{f}} B_{\mathbf{f}}^{\kappa-1}. \quad (8)$$

(a) \Rightarrow (b). If (a) holds $\{L_{\gamma}\mathbf{f}\}_{\gamma \in \Gamma}$ is a Bessel system; thus, having in mind Theorem 5, the entries of the matrix-valued function $\mathbf{F}(\xi)$ belong to $L^{\infty}(\widehat{N})$. Using the Rayleigh-Ritz theorem [14] and Lemma 3, for any $\mathbf{a} \in \ell^2(N)$ we obtain that

$$\begin{aligned} \left\| \sum_{\gamma \in \Gamma} \mathbf{a}(\gamma) L_{\gamma} \mathbf{f} \right\|_{\ell^2(\Gamma)}^2 &= \int_{\widehat{N}} [\mathcal{T}_{\Gamma} \mathbf{a}(\xi)]^* \mathbf{F}^*(\xi) \mathbf{F}(\xi) \mathcal{T}_{\Gamma} \mathbf{a}(\xi) d\mu_{\widehat{N}}(\xi) \\ &\geq \int_{\widehat{N}} \lambda_{\min}[\mathbf{F}^*(\xi) \mathbf{F}(\xi)] \|\mathcal{T}_{\Gamma} \mathbf{a}(\xi)\|^2 d\xi \\ &\geq A_{\mathbf{f}} \int_{\widehat{N}} \|\mathcal{T}_{\Gamma} \mathbf{a}(\xi)\|^2 d\xi = A_{\mathbf{f}} \|\mathcal{T}_{\Gamma} \mathbf{a}\|_{L_{\kappa}^2(\widehat{N})}^2 = A_{\mathbf{f}} \|\mathbf{a}\|_{\ell^2(\Gamma)}^2. \end{aligned} \quad (9)$$

For any number $J > A_{\mathbf{f}}$, there exist a subset $\Omega \subset \widehat{N}$ with positive measure such that $\lambda_{\min}[\mathbf{F}^*(\xi)\mathbf{F}(\xi)] < J$ for $\xi \in \Omega$. Let $\mathbf{a} \in \ell^2(\Gamma)$ such that $\mathcal{T}_{\Gamma} \mathbf{a}(\xi)$ is equal to 0 when $\xi \notin \Omega$, and it is equal to a unitary eigenvector of $\mathbf{F}^*(\xi)\mathbf{F}(\xi)$ corresponding to the eigenvalue $\lambda_{\min}[\mathbf{F}^*(\xi)\mathbf{F}(\xi)]$ when $\xi \in \Omega$. Then, using (6), we obtain

$$\left\| \sum_{\gamma \in \Gamma} \mathbf{a}(\gamma) L_{\gamma} \mathbf{f}(\gamma) \right\|_{\ell^2(\Gamma)}^2 = \int_{\widehat{N}} \lambda_{\min}[\mathbf{F}^*(\xi)\mathbf{F}(\xi)] \|\mathcal{T}_{\Gamma} \mathbf{a}(\xi)\|^2 d\xi \leq J \int_{\widehat{N}} \|\mathcal{T}_{\Gamma} \mathbf{a}(\xi)\|^2 d\xi = J \|\mathbf{a}\|_{\ell^2(\Gamma)}^2.$$

Therefore, if $\{L_{\gamma}\mathbf{f}\}_{\gamma \in \Gamma}$ is a Riesz basis then $A_{\mathbf{f}} > 0$ which, having in mind (8), implies $\operatorname{ess\,inf}_{\xi \in \widehat{N}} |\det \mathbf{F}(\xi)| > 0$. Besides, the optimal lower Riesz bound is $A_{\mathbf{f}}$.

(b) \Rightarrow (a). Since the entries of $\mathbf{F}(\xi) \in L^{\infty}(\widehat{N})$, using Theorem 5 we deduce that $B_{\mathbf{f}} < \infty$. Since $\operatorname{ess\,inf}_{\xi \in \widehat{N}} |\det \mathbf{F}(\xi)| > 0$, using (8) we obtain $A_{\mathbf{f}} > 0$. As a consequence of Theorem 5 and inequality (9) we deduce that the system $\{L_{\gamma}\mathbf{f}\}_{\gamma \in \Gamma}$ is a Riesz basis for $\ell^2(\Gamma)$.

Next, we find its dual Riesz basis. Since $\operatorname{ess\,inf}_{\xi \in \widehat{N}} |\det \mathbf{F}(\xi)| > 0$, and the entries of $\mathbf{F}^*(\xi)$ belongs to $L^{\infty}(\widehat{N})$, the entries of the matrix-valued function $[\mathbf{F}^*(\xi)]^{-1}$ belong to $\ell^2(\widehat{N})$. By Lemma 3, there exist a unique $\mathbf{g} \in \ell^2(\Gamma)$ such that $\mathcal{T}_{\Gamma} \mathbf{g}$ is the first column of $[\mathbf{F}^*(\xi)]^{-1}$, or equivalently, a unique $\mathbf{g} \in \ell^2(\Gamma)$ satisfying (7). From Theorem 4 we get

$$\mathcal{T}_{\Gamma} [\langle \mathbf{g}, L_{\gamma} \mathbf{f} \rangle]_{\gamma \in \Gamma}(\xi) = \mathcal{T}_{\Gamma}(\mathcal{A}_{\mathbf{f}} \mathbf{g})(\xi) = \mathbf{F}^*(\xi) \mathcal{T}_{\Gamma} \mathbf{g}(\xi) = [1 \ 0 \ \dots \ 0]^{\top}$$

Hence $\langle \mathbf{g}, L_{\gamma} \mathbf{f} \rangle = \delta(\gamma)$, and the system $\{L_{\gamma} \mathbf{g}\}_{\gamma \in \Gamma}$ is the dual Riesz basis to $\{L_{\gamma} \mathbf{f}\}_{\gamma \in \Gamma}$. Besides, we have that $\mathbf{g} * \mathbf{f}^*(\gamma) = \langle \mathbf{g}, L_{\gamma} \mathbf{f} \rangle = \delta(\gamma)$. Applying (5), having in mind that the matrices corresponding to δ and \mathbf{f}^* are $[\widehat{\delta}_{h,l}]_{h,l \in H} = \mathbf{I}_{\kappa}$ and $[\widehat{\mathbf{f}}_{h,l}^*]_{h,l \in H} = \mathbf{F}^*$ we get that $\mathbf{F}^*(\xi) \mathbf{G}(\xi) = \mathbf{I}_{\kappa}$, a.e. $\xi \in \widehat{N}$. \square

Corollary 7. *The system $\{L_\gamma \mathbf{f}\}_{\gamma \in \Gamma}$ is an orthonormal basis for $\ell^2(\Gamma)$ if and only if the matrix-valued function $\mathbf{F}(\xi)$ is unitary a.e. $\xi \in \widehat{N}$, or equivalently, if $\mathbf{F}^*(\xi)\mathcal{T}_\Gamma \mathbf{f}(\xi) = [1 \ 0 \ \dots \ 0]^\top$ a.e. $\xi \in \widehat{N}$.*

Proof. The Riesz basis $\{L_\gamma \mathbf{f}\}_{\gamma \in \Gamma}$ is an orthonormal basis if and only if the generator of the dual Riesz basis $\mathbf{g} = \mathbf{f}$, or equivalently, if $\mathbf{G} = \mathbf{F}$. Thus, the result follows from Theorem 6. \square

Corollary 8. *Consider $\mathbf{f}, \mathbf{g} \in \ell^2(\Gamma)$ and their associated transfer matrices $\mathbf{F}(\xi), \mathbf{G}(\xi)$ defined in (3). Assume that the entries of $\mathbf{F}(\xi)$ and $\mathbf{G}(\xi)$ belong to $L^\infty(\widehat{N})$. Then, the systems $\{L_\gamma \mathbf{f}\}_{\gamma \in \Gamma}$ and $\{L_\gamma \mathbf{g}\}_{\gamma \in \Gamma}$ form a pair of dual Riesz bases if and only if $\mathbf{G}(\xi) = [\mathbf{F}^*(\xi)]^{-1}$ a.e. $\xi \in \widehat{N}$.*

Proof. Since the entries of $\mathbf{G}(\xi)$ belong to $L^\infty(\widehat{N})$, if $\mathbf{G}(\xi)\mathbf{F}^*(\xi) = \mathbf{I}_\kappa$ we have that

$$\operatorname{ess\,inf}_{\xi \in \widehat{N}} |\det \mathbf{F}(\xi)| = \operatorname{ess\,inf}_{\xi \in \widehat{N}} (|\det \mathbf{G}(\xi)|^{-1}) = \left(\operatorname{ess\,sup}_{\xi \in \widehat{N}} |\det \mathbf{G}(\xi)| \right)^{-1} > 0.$$

Hence, the result is easily obtained from Theorem 6. \square

3.1 Remarks

- Whenever the generator \mathbf{f} belongs to $\ell^1(\Gamma)$, the matrix-valued function $\mathbf{F}(\xi)$ has continuous entries in the compact \widehat{N} (recall that N is discrete). Thus, from Theorem 5 the system $\{L_\gamma \mathbf{f}\}_{\gamma \in \Gamma}$ is always a Bessel sequence for $\ell^2(\Gamma)$. From Theorem 6 it is a Riesz basis for $\ell^2(\Gamma)$ if and only if the matrix-valued function $\mathbf{F}(\xi)$ is non-singular for all $\xi \in \widehat{N}$. Finally, from Corollary 7 it is an orthonormal basis if and only if $\mathbf{F}(\xi)$ is unitary for all $\xi \in \widehat{N}$.

- Theorems 5 and 6, and Corollary 7 can be restated in terms of the convolution operator. Namely (see [8, Lemma 3.2.1 and Proposition 3.6.8]),

- a. The expression $\Lambda_f \mathbf{a} = \sum_{\eta \in \Gamma} \mathbf{a}(\eta) L_\eta \mathbf{f} = \mathbf{a} * \mathbf{f}$ defines a bounded linear operator $\Lambda_f : \ell^2(\Gamma) \rightarrow \ell^2(\Gamma)$ if and only if the entries of the matrix-valued function $\mathbf{F}(\xi)$ belong to $L^\infty(\widehat{N})$. In this case, $\|\Lambda_f\| = B_f^{1/2}$.

- b. Assume that the entries of the matrix-valued function $\mathbf{F}(\xi)$ belong to $L^\infty(\widehat{N})$. Then, Λ_f is an invertible operator if and only if $\operatorname{ess\,inf}_{\xi \in \widehat{N}} |\det \mathbf{F}(\xi)| > 0$. In this case, $\|\Lambda_f^{-1}\| = A_f^{-1/2}$ and the condition number of Λ_f is $\|\Lambda_f\| \|\Lambda_f^{-1}\| = (B_f/A_f)^{1/2}$. Besides, there exists a unique $\mathbf{h} \in \ell^2(\Gamma)$ such that $\mathbf{f} * \mathbf{h} = \boldsymbol{\delta}$, and satisfying

$$\mathbf{H}(\xi)\mathbf{F}(\xi) = \mathbf{I}_\kappa, \quad \text{a.e. } \xi \in \widehat{N},$$

where $\mathbf{H}(\xi)$ is the transfer matrix associated to \mathbf{h} . Besides, $\mathbf{h} = \mathbf{g}^*$ where \mathbf{g} is the function defined in Theorem 6.

- c. The linear map Λ_f defines a unitary operator if and only if $\mathbf{F}(\xi)$ is unitary a.e. $\xi \in \widehat{N}$.

3.2 Riesz bases examples associated to the infinite dihedral group D_∞

We illustrate the above results of this section in the case of the infinite dihedral group $D_\infty = \mathbb{Z} \rtimes \{1, -1\}$. Recall that $\widehat{\mathbb{Z}} \cong \mathbb{T}$, with $\langle n, z \rangle = z^n$, $z \in \mathbb{T}$, and consequently the Fourier transform of the sequence $\{\mathbf{a}(n)\}_{n \in \mathbb{Z}}$ is the z -transform $\widehat{\mathbf{a}}(z) = \sum_{n \in \mathbb{Z}} \mathbf{a}(n)z^{-n}$ (see, for instance, [10, Theorem 4.5]).

For any $\mathbf{f} \in \ell^2(D_\infty)$, the first column of $\mathbf{F}(z)$, is formed by the z -transforms of $\mathbf{f}_1(n) = \mathbf{f}(n, 1)$ and $\mathbf{f}_{-1}(n) = \mathbf{f}(n, -1)$, $n \in \mathbb{Z}$. The second column is formed by the z -transforms of $\mathbf{f}_{1,-1}(n) = \mathbf{f}_{-1}(-n)$ and $\mathbf{f}_{-1,-1}(n) = \mathbf{f}_1(-n)$, $n \in \mathbb{Z}$; that is

$$\mathbf{F}(z) = \begin{bmatrix} \widehat{\mathbf{f}}_1(z) & \widehat{\mathbf{f}}_{-1}(z^{-1}) \\ \widehat{\mathbf{f}}_{-1}(z) & \widehat{\mathbf{f}}_1(z^{-1}) \end{bmatrix}, \quad z \in \mathbb{T}. \quad (10)$$

Firstly, according to Corollary 7, the sequence $\{L_\gamma \mathbf{f}\}_{\gamma \in D_\infty}$ is an orthonormal basis for $\ell^2(D_\infty)$ if and only if $\mathbf{F}^*(z)\mathcal{T}_\Gamma \mathbf{f}(z) = [1 \ 0]^\top$, or equivalently

$$|\widehat{\mathbf{f}}_1(z)|^2 + |\widehat{\mathbf{f}}_{-1}(z)|^2 = 1 \quad \text{and} \quad \widehat{\mathbf{f}}_1(z)\overline{\widehat{\mathbf{f}}_{-1}(z^{-1})} + \widehat{\mathbf{f}}_{-1}(z)\overline{\widehat{\mathbf{f}}_1(z^{-1})} = 0, \quad \text{a.e. } z \in \mathbb{T}.$$

These equations are satisfied, for example, when $\widehat{\mathbf{f}}_{-1}(z) = 0$ and $|\widehat{\mathbf{f}}_1(z)| = 1$. In signal processing jargon, complex transfer functions satisfying $|f(z)| = 1$ in \mathbb{T} are called allpass filters; expressions for rational allpass filter, their properties, as well as efficient ways to compute the $*_N$ -convolutions in (1) and (4), can be found in Ref. [26, Section 3.4]. The simplest allpass filter $\widehat{\mathbf{f}}_1(z) = z^k$ yields to the canonical basis $\{L_\gamma \boldsymbol{\delta}\}_{\gamma \in D_\infty}$. Other interesting solutions of the above equations are

$$\widehat{\mathbf{f}}_1(e^{iw}) = \begin{cases} 1, & |w| \leq a \\ 0, & |w| > a \end{cases}; \quad \widehat{\mathbf{f}}_{-1}(e^{iw}) = \begin{cases} 0, & |w| \leq a \\ 1, & |w| > a \end{cases}$$

for a fixed $a \in (0, \pi)$.

Secondly, according to Theorem 6, the sequence $\{L_\gamma \mathbf{f}\}_{\gamma \in D_\infty}$ is a Riesz basis for $\ell^2(D_\infty)$ if and only if

$$\operatorname{ess\,inf}_{z \in \mathbb{T}} |\det \mathbf{F}(z)| = \operatorname{ess\,inf}_{z \in \mathbb{T}} |\widehat{\mathbf{f}}_1(z)\widehat{\mathbf{f}}_1(z^{-1}) - \widehat{\mathbf{f}}_{-1}(z)\widehat{\mathbf{f}}_{-1}(z^{-1})| > 0.$$

In this case, by solving $\mathbf{F}^*(z) [\widehat{\mathbf{g}}_1(z) \ \widehat{\mathbf{g}}_{-1}(z)]^\top = [1 \ 0]^\top$, we obtain

$$\widehat{\mathbf{g}}_1(z) = \frac{\overline{\widehat{\mathbf{f}}_1(z)}}{\det \mathbf{F}^*(z)}; \quad \widehat{\mathbf{g}}_{-1}(z) = \frac{-\overline{\widehat{\mathbf{f}}_{-1}(z)}}{\det \mathbf{F}^*(z)}, \quad (11)$$

which provides the generator \mathbf{g} of its dual Riesz basis $\{L_\gamma \mathbf{g}\}_{\gamma \in D_\infty}$. Whether \mathbf{f} is real, we have that $\widehat{\mathbf{f}}_1(z^{-1}) = \overline{\widehat{\mathbf{f}}_1(z)}$ and $\widehat{\mathbf{f}}_{-1}(z^{-1}) = \overline{\widehat{\mathbf{f}}_{-1}(z)}$, from which it is straightforward to deduce that the optimal Riesz bounds are

$$\begin{aligned} A_{\mathbf{f}} &:= \operatorname{ess\,inf}_{z \in \mathbb{T}} \lambda_{\min}[\mathbf{F}^*(z)\mathbf{F}(z)] = \operatorname{ess\,inf}_{w \in [-\pi, \pi]} (|\widehat{\mathbf{f}}_1(e^{iw})| - |\widehat{\mathbf{f}}_{-1}(e^{iw})|)^2, \\ B_{\mathbf{f}} &:= \operatorname{ess\,sup}_{z \in \mathbb{T}} \lambda_{\max}[\mathbf{F}^*(z)\mathbf{F}(z)] = \operatorname{ess\,sup}_{w \in [-\pi, \pi]} (|\widehat{\mathbf{f}}_1(e^{iw})| + |\widehat{\mathbf{f}}_{-1}(e^{iw})|)^2. \end{aligned} \quad (12)$$

Closing this section we exhibit a simple example. The generators \mathbf{f} and \mathbf{g} are finitely supported whenever $\widehat{\mathbf{f}}_1$ and $\widehat{\mathbf{f}}_{-1}$ are Laurent polynomials such that $\det \mathbf{F}(z) = \widehat{\mathbf{f}}_1(z)\widehat{\mathbf{f}}_1(z^{-1}) - \widehat{\mathbf{f}}_{-1}(z)\widehat{\mathbf{f}}_{-1}(z^{-1}) = az^k$ for some $k \in \mathbb{Z}$, $a \neq 0$. For instance, for $\widehat{\mathbf{f}}_1(z) = 3/8$ and $\widehat{\mathbf{f}}_{-1}(z) = z/8$ we obtain $\widehat{\mathbf{g}}_1(z) = 3$ and $\widehat{\mathbf{g}}_{-1}(z) = -z$; thus the dual generators \mathbf{f} and \mathbf{g} have both support of size 2. From (12), the optimal Riesz bounds of $\{L_\gamma \mathbf{f}\}_{\gamma \in \Gamma}$ are $A_{\mathbf{f}} = 1/4$ and $B_{\mathbf{f}} = 1/2$.

4 A sampling application

Let $\Gamma \ni \gamma \mapsto U(\gamma) \in \mathcal{U}(\mathcal{H})$ be a unitary representation of the group Γ in a separable Hilbert space \mathcal{H} , i.e., a homomorphism between Γ and $\mathcal{U}(\mathcal{H})$. We are interested in the study of sampling in the principal U -invariant space $\mathcal{A}_\varphi := \overline{\text{span}}\{U(\gamma)\varphi\}_{\gamma \in \Gamma}$ of \mathcal{H} , where φ denotes a fixed element of \mathcal{H} . In case the sequence $\{U(\gamma)\varphi\}_{\gamma \in \Gamma}$ is a Riesz sequence for \mathcal{H} (one can find necessary and sufficient conditions in Refs. [2, 21]) the subspace \mathcal{A}_φ can be expressed as

$$\mathcal{A}_\varphi = \left\{ \sum_{\gamma \in \Gamma} \mathbf{a}(\gamma)U(\gamma)\varphi : \mathbf{a} = \{\mathbf{a}(\gamma)\}_{\gamma \in \Gamma} \in \ell^2(\Gamma) \right\}$$

The goal here is the stable recovery of any $f \in \mathcal{A}_\varphi$ from the data $\{\mathcal{L}_\psi f(\gamma)\}_{\gamma \in \Gamma}$ given by

$$\mathcal{L}_\psi f(\gamma) := \langle f, U(\gamma)\psi \rangle_{\mathcal{H}}, \quad \gamma \in \Gamma, \quad (13)$$

where $\psi \in \mathcal{H}$ is a fixed element which does not belong necessarily to \mathcal{A}_φ . First, we express the samples in a more suitable manner; for each $f = \sum_{\gamma \in \Gamma} \mathbf{a}(\gamma)U(\gamma)\varphi$ in \mathcal{A}_φ we have

$$\begin{aligned} \mathcal{L}_\psi f(\gamma) &= \left\langle \sum_{\eta \in \Gamma} \mathbf{a}(\eta)U(\eta)\varphi, U(\gamma)\psi \right\rangle_{\mathcal{H}} = \sum_{\eta \in \Gamma} \mathbf{a}(\eta) \langle \varphi, U(\eta^{-1}\gamma)\psi \rangle_{\mathcal{H}} \\ &= \sum_{\eta \in \Gamma} \mathbf{a}(\eta) \overline{\langle \varphi, U(\eta^{-1}\gamma)\psi \rangle_{\mathcal{H}}} = \sum_{\eta \in \Gamma} \mathbf{a}(\eta) \overline{L_\eta \mathbf{f}_\psi} = \langle \mathbf{a}, L_\gamma \mathbf{f}_\psi \rangle_{\ell^2(\Gamma)}, \quad \gamma \in \Gamma, \end{aligned}$$

where $\mathbf{f}_\psi(\eta) := \overline{\langle \varphi, U(\eta^{-1})\psi \rangle_{\mathcal{H}}}$ for $\eta \in \Gamma$, and $\mathbf{a} = \{\mathbf{a}(\gamma)\}_{\gamma \in \Gamma}$. Notice that \mathbf{f}_ψ belongs to $\ell^2(\Gamma)$. In the light of Theorem 6, assume that $\{L_\gamma \mathbf{f}_\psi\}_{\gamma \in \Gamma}$ is a Riesz basis for $\ell^2(\Gamma)$ with dual Riesz basis $\{L_\gamma \mathbf{g}_\psi\}_{\gamma \in \Gamma}$. Thus, for any $\mathbf{a} \in \ell^2(\Gamma)$ we have

$$\mathbf{a} = \sum_{\gamma \in \Gamma} \langle \mathbf{a}, L_\gamma \mathbf{f}_\psi \rangle_{\ell^2(\Gamma)} L_\gamma \mathbf{g}_\psi = \sum_{\gamma \in \Gamma} \mathcal{L}_\psi f(\gamma) L_\gamma \mathbf{g}_\psi \quad \text{in } \ell^2(\Gamma). \quad (14)$$

In order to derive a sampling formula in \mathcal{A}_φ compatible with its structure, we consider the natural isomorphism $\mathcal{T}_{U,\varphi} : \ell^2(\Gamma) \rightarrow \mathcal{A}_\varphi$ which maps the usual orthonormal basis $\{\delta_\gamma\}_{\gamma \in \Gamma}$ for $\ell^2(\Gamma)$ onto the Riesz basis $\{U(\gamma)\varphi\}_{\gamma \in \Gamma}$ for \mathcal{A}_φ . This isomorphism satisfies the following *shifting property*:

$$\mathcal{T}_{U,\varphi}(L_\gamma \mathbf{f}) = U(\gamma)\mathcal{T}_{U,\varphi} \mathbf{f} \quad \text{for each } \mathbf{f} \in \ell^2(\Gamma) \text{ and } \gamma \in \Gamma.$$

Indeed, we have that $L_\gamma \delta_\eta = \delta_{\gamma\eta}$ for $\gamma, \eta \in \Gamma$. As a consequence, $\mathcal{T}_{U,\varphi}(L_\gamma \delta_\eta) = \mathcal{T}_{U,\varphi} \delta_{\gamma\eta} = U(\gamma)U(\eta)\varphi = U(\gamma)\mathcal{T}_{U,\varphi}(\delta_\eta)$. From a continuity argument the result becomes true for any $\mathbf{f} \in \ell^2(\Gamma)$.

Now, consider $f = \mathcal{T}_{U,\varphi}(\mathbf{a})$ in \mathcal{A}_φ ; applying the isomorphism $\mathcal{T}_{U,\varphi}$ in expansion (14) and using the above shifting property we obtain for each $f \in \mathcal{A}_\varphi$ the sampling formula

$$f = \mathcal{T}_{U,\varphi}(\mathbf{a}) = \sum_{\gamma \in \Gamma} \mathcal{L}_\psi f(\gamma) \mathcal{T}_{U,\varphi}(L_\gamma \mathbf{g}_\psi) = \sum_{\gamma \in \Gamma} \mathcal{L}_\psi f(\gamma) U(\gamma) \mathcal{T}_{U,\varphi}(\mathbf{g}_\psi) \quad \text{in } \mathcal{H}. \quad (15)$$

Notice that $\mathcal{T}_{U,\varphi}(\mathbf{g}_\psi) = \sum_{\gamma \in \Gamma} \mathbf{g}_\psi(\gamma) U(\gamma) \varphi \in \mathcal{A}_\varphi$. Moreover, since $\mathcal{T}_{U,\varphi}$ is an isomorphism, the sequence $\{U(\gamma) \mathcal{T}_{U,\varphi}(\mathbf{g}_\psi)\}_{\gamma \in \Gamma}$ is a Riesz basis for \mathcal{A}_φ . In fact, the following sampling theorem in \mathcal{A}_φ holds:

Theorem 9. *For a given $\psi \in \mathcal{H}$, consider $\mathbf{f}_\psi \in \ell^2(\Gamma)$ such that $\mathbf{f}_\psi(\eta) := \langle \varphi, U(\eta^{-1})\psi \rangle_{\mathcal{H}}$ for $\eta \in \Gamma$. Assume that all the entries of its associated $\kappa \times \kappa$ matrix-valued function $\mathbf{F}(\xi)$ defined in (3) belong to $L^\infty(\widehat{N})$. The following statements are equivalent:*

- (a) $\text{ess inf}_{\xi \in \widehat{N}} |\det \mathbf{F}(\xi)| > 0$.
- (b) *There exists a unique $\mathbf{g}_\psi \in \ell^2(\Gamma)$ such that its associate matrix-valued function $\mathbf{G}(\xi)$ defined in (3) has entries in $L^\infty(\widehat{N})$, and it satisfies $\mathbf{G}(\xi) \mathbf{F}^*(\xi) = \mathbf{I}_\kappa$, a.e. $\xi \in \widehat{N}$.*
- (c) *There exists a unique $\Phi_\psi \in \mathcal{A}_\varphi$ such that the sequence $\{U(\gamma) \Phi_\psi\}_{\gamma \in \Gamma}$ is a Riesz basis for \mathcal{A}_φ and the sampling formula*

$$f = \sum_{\gamma \in \Gamma} \mathcal{L}_\psi f(\gamma) U(\gamma) \Phi_\psi \quad \text{in } \mathcal{H} \quad (16)$$

holds for each $f \in \mathcal{A}_\varphi$.

In case the equivalent conditions are satisfied, necessarily $\Phi_\psi = \mathcal{T}_{U,\varphi}(\mathbf{g}_\psi)$ where $\mathbf{g}_\psi \in \ell^2(\Gamma)$ satisfies conditions in (b). Moreover, the interpolation property $\mathcal{L}_\psi \Phi_\psi(\gamma) = \delta_{\gamma, e_\Gamma}$, $\gamma \in \Gamma$, holds.

Proof. (a) \Rightarrow (b). The sequence $\{L_\gamma \mathbf{f}_\psi\}_{\ell^2(\Gamma)}$ is a Riesz basis for $\ell^2(\Gamma)$. Having in mind Theorem 6, its dual Riesz basis has the form $\{L_\gamma \mathbf{g}_\psi\}_{\ell^2(\Gamma)}$ with $\mathbf{G}(\xi) \mathbf{F}^*(\xi) = \mathbf{I}_\kappa$, a.e. $\xi \in \widehat{N}$.

(b) \Rightarrow (c). According with Corollary 8, the sequences $\{L_\gamma \mathbf{f}_\psi\}_{\ell^2(\Gamma)}$ and $\{L_\gamma \mathbf{g}_\psi\}_{\ell^2(\Gamma)}$ form a pair of dual Riesz bases for $\ell^2(\Gamma)$. Thus we have (14) and, consequently, (15) proves (c).

(c) \Rightarrow (a). Applying the isomorphism $\mathcal{T}_{U,\varphi}^{-1}$, the sequence $\{\mathcal{T}_{U,\varphi}^{-1}(U(\gamma) \Phi_\psi)\}_{\gamma \in \Gamma}$ is a Riesz sequence for $\ell^2(\Gamma)$, and for each $\mathbf{a} \in \ell^2(\Gamma)$ we get

$$\mathbf{a} = \sum_{\gamma \in \Gamma} \langle \mathbf{a}, L_\gamma \mathbf{f}_\psi \rangle_{\ell^2(\Gamma)} \mathcal{T}_{U,\varphi}^{-1}(U(\gamma) \Phi_\psi) \quad \text{in } \ell^2(\Gamma).$$

The sequence $\{L_\gamma \mathbf{f}_\psi\}_{\ell^2(\Gamma)}$ is a Bessel sequence biorthogonal to $\{\mathcal{T}_{U,\varphi}^{-1}(U(\gamma) \Phi_\psi)\}_{\gamma \in \Gamma}$, and hence it is a Riesz basis for $\ell^2(\Gamma)$ [8, Theorem 3.6.7]; from Theorem 6, $\text{ess inf}_{\xi \in \widehat{N}} |\det \mathbf{F}(\xi)| > 0$.

The uniqueness of the coefficients in a Riesz basis expansion gives the interpolation property $\mathcal{L}_\psi \Phi_\psi(\gamma) = \delta_{\gamma, e_\Gamma}$, $\gamma \in \Gamma$. \square

4.1 The crystallographic group case

The *euclidean motion group* $E(d)$ is the semidirect product $\mathbb{R}^d \rtimes_{\sigma} O(d)$ corresponding to the homomorphism $\sigma : O(d) \rightarrow \text{Aut}(\mathbb{R}^d)$ given by $\sigma(A)(x) = Ax$, where $A \in O(d)$ and $x \in \mathbb{R}^d$. The composition law on $E(d) = \mathbb{R}^d \rtimes_{\sigma} O(d)$ reads $(x, A) \cdot (x', A') = (x + Ax', AA')$.

Let \mathcal{M} be a non-singular $d \times d$ matrix and H a finite subgroup of $O(d)$ of order κ such that $A(\mathcal{M}\mathbb{Z}^d) = \mathcal{M}\mathbb{Z}^d$ for each $A \in H$. We consider the *crystallographic group* $\Gamma_{\mathcal{M},H} := \mathcal{M}\mathbb{Z}^d \rtimes_{\sigma} H$ and its *quasi regular representation* (see, for instance, Ref. [2]) on $L^2(\mathbb{R}^d)$

$$U(n, A)f(t) = f[A^{\top}(t - n)], \quad n \in \mathcal{M}\mathbb{Z}^d, A \in H \text{ and } f \in L^2(\mathbb{R}^d).$$

For a fixed $\varphi \in L^2(\mathbb{R}^d)$ such that the sequence $\{U(n, A)\varphi\}_{(n,A) \in \Gamma_{\mathcal{M},H}}$ is a Riesz sequence for $L^2(\mathbb{R}^d)$ (see, for instance, Refs.[7, 15]) we consider the U -invariant subspace in $L^2(\mathbb{R}^d)$

$$\mathcal{A}_{\varphi} = \left\{ \sum_{(n,A) \in \Gamma_{\mathcal{M},H}} \alpha(n, A) \varphi[A^{\top}(t - n)] : \{\alpha(n, A)\} \in \ell^2(\Gamma_{\mathcal{M},H}) \right\}$$

For a fixed $\psi \in L^2(\mathbb{R}^d)$ non necessarily in \mathcal{A}_{φ} we consider the average samples of any $f \in \mathcal{A}_{\varphi}$

$$\mathcal{L}_{\psi}f(n, A) = \langle f, U(n, A)\psi \rangle_{L^2(\mathbb{R}^d)} = \langle f, \psi[A^{\top}(\cdot - n)] \rangle_{L^2(\mathbb{R}^d)}, \quad (n, A) \in \Gamma_{\mathcal{M},H}.$$

Under the hypotheses in Theorem 9, there exists a function $\Phi_{\psi} \in \mathcal{A}_{\varphi}$ such that the sequence $\{\Phi_{\psi}[A^{\top}(t - n)]\}_{(n,A) \in \Gamma_{\mathcal{M},H}}$ is a Riesz basis for \mathcal{A}_{φ} , and for each $f \in \mathcal{A}_{\varphi}$ we have the sampling expansion

$$f(t) = \sum_{(n,A) \in \Gamma_{\mathcal{M},H}} \langle f, \psi[A^{\top}(\cdot - n)] \rangle_{L^2(\mathbb{R}^d)} \Phi_{\psi}[A^{\top}(t - n)] \quad \text{in } L^2(\mathbb{R}^d). \quad (17)$$

If the generator $\varphi \in C(\mathbb{R}^d)$ and the function $t \mapsto \sum_{(n,A) \in \Gamma_{\mathcal{M},H}} |\varphi[A^{\top}(t - n)]|^2$ is bounded on \mathbb{R}^d , a standard argument shows that \mathcal{A}_{φ} is a reproducing kernel Hilbert space (RKHS) of continuous bounded functions in $L^2(\mathbb{R}^d)$. As a consequence, convergence in $L^2(\mathbb{R}^d)$ -norm implies pointwise convergence which is absolute and uniform on \mathbb{R}^d .

4.2 The pointwise samples case

Let $\{U(\gamma)\}_{\gamma \in \Gamma}$ be a unitary representation of the group $\Gamma = N \rtimes_{\sigma} H$ on the Hilbert space $\mathcal{H} = L^2(\mathbb{R}^d)$. If the generator $\varphi \in L^2(\mathbb{R}^d)$ satisfies that, for each $\gamma \in \Gamma$, the function $U(\gamma)\varphi$ is continuous on \mathbb{R}^d , and the condition

$$\sup_{t \in \mathbb{R}^d} \sum_{\gamma \in \Gamma} |[U(\gamma)\varphi](t)|^2 < +\infty, \quad (18)$$

then the subspace \mathcal{A}_{φ} is a RKHS of continuous bounded functions in $L^2(\mathbb{R}^d)$. In fact, the following result holds:

Lemma 10. *For any $\{\mathbf{a}(\gamma)\}_{\gamma \in \Gamma} \in \ell^2(\Gamma)$ the series $\sum_{\gamma \in \Gamma} \mathbf{a}(\gamma) [U(\gamma)\varphi](t)$ converges pointwise to a continuous bounded function if and only if for each $\gamma \in \Gamma$, the function $U(\gamma)\varphi$ is continuous on \mathbb{R}^d , and condition (18) holds.*

Proof. Cauchy-Schwarz inequality and Weierstrass M-test prove the sufficient condition. To prove the necessary condition we follow the arguments in [27]. Indeed, notice first that choosing the delta sequences in $\ell^2(\Gamma)$ we deduce that each function $U(\gamma)\varphi$ is continuous on \mathbb{R}^d .

For each fixed $t \in \mathbb{R}^d$, since the series $\sum_{\gamma \in \Gamma} \mathbf{a}(\gamma) [U(\gamma)\varphi](t)$ converges for any $\{\mathbf{a}(\gamma)\}_{\gamma \in \Gamma}$ in $\ell^2(\Gamma)$, we obtain that $\sum_{\gamma \in \Gamma} |[U(\gamma)\varphi](t)|^2 < +\infty$. Moreover, the functional $\Omega_t : \ell^2(\Gamma) \rightarrow \mathbb{C}$ defined as $\Omega_t \mathbf{a} := \sum_{\gamma \in \Gamma} \mathbf{a}(\gamma) [U(\gamma)\varphi](t)$ is bounded with norm $\|\Omega_t\|^2 = \sum_{\gamma \in \Gamma} |[U(\gamma)\varphi](t)|^2$ (see, for instance, [13, p.145]). Next, for fixed $\mathbf{a} = \{\mathbf{a}(\gamma)\}_{\gamma \in \Gamma} \in \ell^2(\Gamma)$ we consider its associated function $f_{\mathbf{a}}(t) := \sum_{\gamma \in \Gamma} \mathbf{a}(\gamma) [U(\gamma)\varphi](t)$, $t \in \mathbb{R}^d$. Since $f_{\mathbf{a}}$ is bounded on \mathbb{R}^d , we get $\sup_{t \in \mathbb{R}^d} |\Omega_t \mathbf{a}| = \sup_{t \in \mathbb{R}^d} |f_{\mathbf{a}}(t)| < +\infty$. Hence, Banach-Steinhaus theorem concludes that

$$\sup_{t \in \mathbb{R}^d} \|\Omega_t\| = \sup_{t \in \mathbb{R}^d} \left(\sum_{\gamma \in \Gamma} |[U(\gamma)\varphi](t)|^2 \right)^{1/2} < +\infty.$$

□

Now for a fixed point $p \in \mathbb{R}^d$ we consider, for each $f \in \mathcal{A}_\varphi$, the new samples given by

$$\mathcal{L}_p f(\gamma) := [U(\gamma^{-1})f](p), \quad \gamma \in \Gamma. \quad (19)$$

For each $f = \sum_{\eta \in \Gamma} \mathbf{a}(\eta) U(\eta)\varphi$ in \mathcal{A}_φ we get

$$\begin{aligned} \mathcal{L}_p f(\gamma) &= \left[\sum_{\eta \in \Gamma} \mathbf{a}(\eta) U(\gamma^{-1}\eta)\varphi \right](p) = \sum_{\eta \in \Gamma} \mathbf{a}(\eta) [U(\gamma^{-1}\eta)\varphi](p) \\ &= \sum_{\eta \in \Gamma} \mathbf{a}(\eta) \overline{f_p(\gamma^{-1}\eta)} = \langle \mathbf{a}, L_\gamma \mathbf{f}_p \rangle_{\ell^2(\Gamma)}, \quad \gamma \in \Gamma, \end{aligned}$$

where $f_p(\eta) := \overline{[U(\eta)\varphi](p)}$, $\eta \in \Gamma$; notice that \mathbf{f}_p belongs to $\ell^2(\Gamma)$. As a consequence, under the hypotheses in Theorem 9 on this new $\mathbf{f}_p \in \ell^2(\Gamma)$, a sampling formula as (16) holds for the data sequence $\{\mathcal{L}_p f(\gamma)\}_{\gamma \in \Gamma}$.

In the particular case of the quasi regular representation of a crystallographic group $\Gamma_{\mathcal{M}, H} = \mathcal{M}\mathbb{Z}^d \rtimes_\sigma H$, for each $f \in \mathcal{A}_\varphi$ its samples (19) are the pointwise samples

$$\mathcal{L}_p f(n, A) = [U[(n, A)^{-1}]f](p) = [U(-A^\top n, A^\top)f](p) = f(Ap + n), \quad (n, A) \in \Gamma.$$

Thus (under hypotheses in Theorem 9), there exists a unique function $\Phi_p \in \mathcal{A}_\varphi$ such that for each $f \in \mathcal{A}_\varphi$ the sampling formula

$$f(t) = \sum_{(n, A) \in \Gamma} f(Ap + n) \Phi_p[A^\top(t - n)], \quad t \in \mathbb{R}^d \quad (20)$$

holds. The convergence of the series in $L^2(\mathbb{R}^d)$ -norm implies pointwise convergence which is absolute and uniform on \mathbb{R}^d . The interpolating function $\Phi_p = \mathcal{T}_{U, \varphi}(\mathbf{g}_p)$ where \mathbf{g}_p is the generator of the dual Riesz basis (see Theorem 6). Coefficients in the expansion $f = \sum_{\gamma \in \Gamma} \mathbf{a}(\gamma) U(\gamma)\varphi$ can be computed from samples as

$$\mathbf{a} = \sum_{\gamma \in \Gamma} f(Ap + n) L_\gamma \mathbf{g}_p \quad (21)$$

4.3 An example involving the infinite dihedral group D_∞

To illustrate the results in this section we consider group $\Gamma = D_\infty$, a unidimensional crystallographic group, and a real generator $\varphi \in L^2(\mathbb{R})$ supported in the interval $[0, 2]$. Notice that we can check if a system $\{U(\gamma)\varphi(t)\}_{\gamma \in D_\infty} = \{\varphi(t-n)\}_{n \in \mathbb{Z}} \cup \{\varphi(n-t)\}_{n \in \mathbb{Z}}$ is a Riesz basis for $\mathcal{A}_\varphi = \{\sum_{n \in \mathbb{Z}} a(n)\varphi(t-n) + b(n)\varphi(n-t) : a, b \in \ell^2(\mathbb{Z})\}$ by using the Gramian condition (see, for instance, Refs. [7, 15]). For instance, the function $\varphi(t) = (16t^2 - 13)t^2(2-t)^2\chi_{[0,2]}(t)$, $t \in \mathbb{R}$, fulfills these requirements.

The aim here is the recovery of any $f \in \mathcal{A}_\varphi$ from its samples $\{f(n+p)\}_{n \in \mathbb{Z}} \cup \{f(n-p)\}_{n \in \mathbb{Z}}$ with a fixed $p \in (0, 1/2)$. We proceed to check condition (a) in Theorem 9. Indeed, since $\text{supp } \varphi \subseteq [0, 2]$, we obtain $\hat{f}_1(z) = \varphi(p) + \varphi(p+1)z$ and $\hat{f}_{-1}(z) = \varphi(1-p)z^{-1} + \varphi(2-p)z^{-2}$ and then (see Eq.(10))

$$\det \mathbf{F}(z) = C + D(z + z^{-1}), \quad z \in \mathbb{T},$$

where $C = \varphi(p)^2 + \varphi(p+1)^2 - \varphi(1-p)^2 - \varphi(2-p)^2$ and $D = \varphi(p)\varphi(p+1) - \varphi(1-p)\varphi(2-p)$. Since $\det \mathbf{F}(e^{iw}) = C + 2D \cos(w)$, whenever $|C| > 2|D|$ the sampling formula (20) holds. It reads

$$f(t) = \sum_{n \in \mathbb{Z}} \{f(n+p)\Phi_p(t-n) + f(n-p)\Phi_p(n-t)\}, \quad t \in \mathbb{R},$$

where the interpolating function is $\Phi_p(t) = \sum_{n \in \mathbb{Z}} \{\mathbf{g}_1(n)\varphi(t-n) + \mathbf{g}_{-1}(n)\varphi(n-t)\}$, $t \in \mathbb{R}$, with (see Eq. (11))

$$\hat{\mathbf{g}}_1(z) = \frac{\varphi(p) + \varphi(p+1)z}{C + D(z + z^{-1})}, \quad \hat{\mathbf{g}}_{-1}(z) = -\frac{\varphi(1-p)z^{-1} + \varphi(2-p)z^{-2}}{C + D(z + z^{-1})}, \quad z \in \mathbb{T}.$$

Note that, whenever $D = 0$, the interpolating function Φ_p has also compact support. For instance, by choosing the generator $\varphi(t) = (16t^2 - 13)t^2(2-t)^2\chi_{[0,2]}(t)$, $t \in \mathbb{R}$, we obtain that $D = 0$ and $C = \frac{3627}{64}$, and therefore

$$\Phi_p(t) = \frac{64}{3627} [\varphi(p)\varphi(t) + \varphi(p+1)\varphi(t+1) - \varphi(1-p)\varphi(1-t) - \varphi(2-p)\varphi(2-t)], \quad t \in \mathbb{R},$$

has support $[-1, 2]$. Using Eqs. (12), the computation of coefficients $\mathbf{a}(\gamma)$ in (21) has condition number $(B_f/A_f)^{1/2} \approx 4.82$. For this choice of φ , the D_∞ -invariant space \mathcal{A}_φ is a subspace of the space of cardinal splines of degree 6 with nodes at \mathbb{Z} and continuous derivative.

5 A C^* -algebra connection

It is known that the Banach space $\ell^1(\Gamma)$ becomes a Banach $*$ -algebra under convolution but it is not a C^* -algebra. To avoid this drawback, it can be used the group C^* -algebra of Γ denoted by $C^*(\Gamma)$; it is the completion of $\ell^1(\Gamma)$ with respect to the norm $\|f\| = \|\Lambda_f\|_{\mathcal{B}(\ell^2(\Gamma))}$ [6, II.10.2].

In Ref. [25] it is proved that the mapping $\mathbf{f} \mapsto \mathbf{F}$ in Definition 1 is a C^* -isomorphism between $C^*(\Gamma)$ and a C^* -subalgebra of $\mathcal{M}_\kappa(C(\hat{N}))$, the C^* -algebra of the $\kappa \times \kappa$ matrices with continuous entries on \hat{N} (see also [22]). Thus, this C^* -isomorphism provides an explicit description for the group C^* -algebra of the semidirect product group Γ .

Next, we show that for the semidirect product group $\Gamma = N \rtimes_\sigma H$, when N is a discrete abelian group and H a finite group, an alternative to the group C^* -algebra of Γ is the larger space

$$L^*(\Gamma) := \left\{ \mathbf{f} \in \ell^2(\Gamma) : \hat{f}_{h,l} \in L^\infty(\hat{N}), h, l \in H \right\}.$$

In Theorem 12 below we will prove that $L^*(\Gamma)$ is a C^* -algebra, and that the linear map \mathcal{S} (see Definition 1)

$$L^*(\Gamma) \ni f \xrightarrow{\mathcal{S}} \mathbf{F} \in \mathcal{M}_\kappa(L^\infty(\widehat{N}))$$

defines a C^* -isomorphism, and consequently an isometry, between the C^* -algebra $L^*(\Gamma)$ and a C^* -subalgebra of $\mathcal{M}_\kappa(L^\infty(\widehat{N}))$. Thus, the space $L^*(\Gamma)$ allows to consider, in a C^* -algebra setting, elements of $\ell^2(\Gamma)$ with discontinuous Fourier transform, such as ideal filters in signal processing applications. In case the group $\Gamma = \mathbb{Z} \rtimes_\sigma 1_H \cong \mathbb{Z}$, the space $L^*(\Gamma)$ coincides with the space $A'(\mathbb{Z})$ of pseudomeasures [4, 3.1.8].

Specifically, by $\mathcal{M}_\kappa(L^\infty(\widehat{N}))$ we denote the involution algebra formed by the $\kappa \times \kappa$ matrices with entries in $L^\infty(\widehat{N})$, with pointwise addition and multiplication and where the involution is given by the adjoint matrix. Any $\mathbf{A} \in \mathcal{M}_\kappa(L^\infty(\widehat{N}))$ can be represented by the bounded operator $\pi_{\mathbf{A}} \in \mathcal{B}(L^2_\kappa(\widehat{N}))$ defined by $(\pi_{\mathbf{A}} \mathbf{x})(\xi) := \mathbf{A}(\xi) \mathbf{x}(\xi)$. With respect to the norm induced by this representation,

$$\|\mathbf{A}\|_{\mathcal{M}_\kappa(L^\infty(\widehat{N}))} := \|\pi_{\mathbf{A}}\|_{\mathcal{B}(L^2_\kappa(\widehat{N}))} = \sup \left\{ \|\mathbf{A}(\cdot) \mathbf{x}(\cdot)\|_{L^2_\kappa(\widehat{N})} : \|\mathbf{x}\|_{L^2_\kappa(\widehat{N})} = 1 \right\},$$

the involution algebra $\mathcal{M}_\kappa(L^\infty(\widehat{N}))$ is a C^* -algebra [6, II.6.6]. An estimation for this norm can be found in Ref.[5].

Lemma 11. *For each $\mathbf{A} \in \mathcal{M}_\kappa(L^\infty(\widehat{N}))$ we have that $\|\mathbf{A}\|_{\mathcal{M}_\kappa(L^\infty(\widehat{N}))} = \text{ess sup}_{\xi \in \widehat{N}} \|\mathbf{A}(\xi)\|_2$, where $\|\mathbf{A}(\xi)\|_2$ denotes the spectral norm of the matrix $\mathbf{A}(\xi)$.*

Proof. Since $\|\pi_{\mathbf{A}} \mathbf{x}\|_{L^2_\kappa(\widehat{N})}^2 = \int_{\widehat{N}} \mathbf{x}^*(\xi) \mathbf{A}^*(\xi) \mathbf{A}(\xi) \mathbf{x}(\xi) d\mu_{\widehat{N}}(\xi)$, the lemma can be proved by means of the argument used to prove Theorem 5 from equality (6). \square

Alternatively, this lemma could be proved by checking that $\mathcal{M}_\kappa(L^\infty(\widehat{N}))$ with the norm $\|\mathbf{A}\| = \text{ess sup}_{\xi \in \widehat{N}} \|\mathbf{A}(\xi)\|_2$ is a C^* -algebra, and having in mind the uniqueness of the C^* -norm.

Theorem 12. *The vector space $L^*(\Gamma)$ under the convolution product, the involution defined by $f^*(\gamma) = \overline{f(-\gamma)}$, $\gamma \in \Gamma$, and the norm $\|f\|_{L^*(\Gamma)} = \|\mathbf{F}\|_{\mathcal{M}_\kappa(L^\infty(\widehat{N}))}$ becomes a C^* -algebra. Besides, the linear map $\mathcal{S} : f \mapsto \mathbf{F}$ is a C^* -isomorphism between $L^*(\Gamma)$ and a C^* -subalgebra of $\mathcal{M}_\kappa(L^\infty(\widehat{N}))$. The transform \mathcal{S} changes the order of the multiplication, i.e., $\mathcal{S}(\mathbf{g} * \mathbf{f}) = \mathcal{S}(\mathbf{f})\mathcal{S}(\mathbf{g})$, $\mathbf{f}, \mathbf{g} \in L^*(\Gamma)$.*

Proof. We can easily check that \mathcal{S} satisfies $\mathcal{S}(f^*) = (\mathcal{S}f)^*$. According to Theorem 4 we have that for any $\mathbf{f}, \mathbf{g} \in L^*(\Gamma)$, $\mathcal{S}(\mathbf{g} * \mathbf{f})(\xi) = \mathcal{S}\mathbf{f}(\xi)\mathcal{S}\mathbf{g}(\xi)$ a.e. $\xi \in \widehat{N}$. As $\|f\|_{L^*(\Gamma)} = \|\mathbf{F}\|_{\mathcal{M}_\kappa(L^\infty(\widehat{N}))} = \|\mathcal{S}\mathbf{f}\|_{\mathcal{M}_\kappa(L^\infty(\widehat{N}))}$, we just need to prove that the range of \mathcal{S} is closed in norm.

In so doing, let us consider $\mathbf{F}_i = \mathcal{S}\mathbf{f}_i$, with $\mathbf{f}_i \in L^*(\Gamma)$, and a matrix $\mathbf{A} \in \mathcal{M}_\kappa(L^\infty(\widehat{N}))$ such that $\|\mathbf{F}_i - \mathbf{A}\|_{\mathcal{M}_\kappa(L^\infty(\widehat{N}))} \rightarrow 0$ as $i \mapsto \infty$. We have to prove that \mathbf{A} belongs to the range of \mathcal{S} . From Lemma 11, $\text{ess sup}_{\xi \in \widehat{N}} \|\mathbf{F}_i(\xi) - \mathbf{A}(\xi)\|_2 \rightarrow 0$ as $i \mapsto \infty$. Having in mind that $\max_{h,l} |b_{h,l}| \leq \|\mathbf{B}\|_2$ for any matrix $\mathbf{B} = [b_{h,l}]$, we obtain that $\text{ess sup}_{\xi \in \widehat{N}} |(\mathbf{F}_i)_{h,l}(\xi) - \mathbf{A}_{h,l}(\xi)| \rightarrow 0$ and then $\|(\mathbf{F}_i)_{h,l} - \mathbf{A}_{h,l}\|_{L^\infty(\widehat{N})} \rightarrow 0$ as $i \mapsto \infty$. Having in mind that \widehat{N} is compact, we also have that

$$\|(\mathbf{F}_i)_{h,l} - \mathbf{A}_{h,l}\|_{L^2(\widehat{N})} \rightarrow 0 \quad \text{as } i \mapsto \infty, \quad l, h \in H. \quad (22)$$

On the other hand, since, for each $h \in H$, $\mathbf{A}_{h,1_H} \in L^\infty(\widehat{N}) \subset L^2(\widehat{N})$, there exists a unique $f \in \ell^2(\Gamma)$ such that the Fourier transform of $f(\cdot, h)$ is $\mathbf{A}_{h,1_H}$.

For any $h \in H$, the sequence $f_i(\cdot, h)$ converges in $\ell^2(N)$ to $f(\cdot, h)$ since, by (22), its Fourier transform $(\mathbf{F}_i)_{h,1}$ converges in $L^2(\widehat{N})$ to $\mathbf{A}_{h,1_H}$ the Fourier transform of $f(\cdot, h)$. Hence, for any $h, l \in H$, the sequence $(f_i)_{h,l} = f_i(-\sigma_l(\cdot), l^{-1}h)$ converges in $\ell^2(N)$ to $f(-\sigma_l(\cdot), l^{-1}h) = f_{h,l}$ and then its Fourier transform $(\mathbf{F}_i)_{h,l}$ converges in $L^2(\widehat{N})$ to $\widehat{f}_{h,l}$. Thus, by using (22) and the uniqueness of the limit we obtain that $\widehat{f}_{h,l} = \mathbf{A}_{h,l}$, $h, l \in H$, and then $\mathcal{S}f = \mathbf{A}$. \square

Theorem 12 gives a simple description of the convolution C^* -algebra $L^*(\Gamma)$. For example, from (10), the C^* -algebra $L^*(D_\infty)$ for the infinite dihedral group D_∞ is C^* -isomorphic to the C^* -algebra of matrices of the type

$$\mathbf{A}(z) = \begin{bmatrix} f(z) & g(z^{-1}) \\ g(z) & f(z^{-1}) \end{bmatrix}, \quad z \in \mathbb{T}, \quad \text{with } f, g \in L^\infty(\mathbb{T}),$$

and the norm $\|\mathbf{A}\|_{\mathcal{M}_2(L^\infty(\mathbb{T}))} = \text{ess sup}_{z \in \mathbb{T}} \|\mathbf{A}(z)\|_2$.

Finally, it is worth to mention that it is possible to give an alternative proof of Theorem 6 by using Theorem 12 and C^* -algebras techniques.

A new proof of Theorem 6:

Proof. Let $\mathcal{B}(\ell^2(\Gamma))$ be the C^* -algebra of bounded linear operators on $\ell^2(\Gamma)$. The linear map

$$L^*(\Gamma) \ni f \xrightarrow{\Lambda} \Lambda_f \in \mathcal{B}(\ell^2(\Gamma))$$

defines a C^* -isomorphism between $L^*(\Gamma)$ and a C^* -subalgebra of $\mathcal{B}(\ell^2(\Gamma))$. Indeed, from Theorem 5 (see the second remark in 3.1), we obtain that any $f \in L^*(\Gamma)$ satisfies $\Lambda_f \in \mathcal{B}(\ell^2(\Gamma))$ and $\|f\|_{L^*(\Gamma)} = \|\mathbf{F}\|_{\mathcal{M}_n(L^\infty(\Gamma))} = \text{ess sup}_{\xi \in \widehat{N}} \|\mathbf{F}(\xi)\|_2 = B_f^{1/2} = \|\Lambda_f\|_{\mathcal{B}(\ell^2(\Gamma))}$; using (5) in Theorem 4 we obtain that $[\Lambda_{f*g}]\mathbf{h} = \Lambda_g(\Lambda_f\mathbf{h})$, for all $f, g \in L^*(\Gamma)$ and $\mathbf{h} \in \ell^2(\Gamma)$; and from (4) we have $\Lambda_{f^*} = \Lambda_f^*$ for $f \in L^*(\Gamma)$.

Hence, from Theorem 12 we deduce that the operator

$$\Lambda(L^*(\Gamma)) \ni \Lambda_f \xrightarrow{\mathcal{S}\Lambda^{-1}} \mathbf{F} \in \mathcal{S}(L^*(\Gamma))$$

is a C^* -isomorphism.

Assume first (a), that is $\{L_\gamma f\}_{\gamma \in \Gamma}$ is a Riesz basis for $\ell^2(\Gamma)$. Then, from Theorem 5, the entries of the matrix-valued function $\mathbf{F}(\xi)$ belong to $L^\infty(\widehat{N})$ and the upper Riesz bound is $B_f < \infty$. Besides, $\Lambda_f \in \mathcal{B}(\ell^2(\Gamma))$ and it is invertible. Since $\Lambda(L^*(\Gamma))$ is a unital C^* -subalgebra of $\mathcal{B}(\ell^2(\Gamma))$, and Λ_f belongs to $\Lambda(L^*(\Gamma))$, its inverse Λ_f^{-1} also belongs to $\Lambda(L^*(\Gamma))$ [24, Proposition 4.8]. Then, by applying the C^* -isomorphism $\mathcal{S}\Lambda^{-1}$, we obtain that \mathbf{F} is invertible in the C^* -subalgebra $\mathcal{S}(L^*(\Gamma))$ and that the lower Riesz bound is [8, Proposition 3.6.8]

$$\begin{aligned} \|\Lambda_f^{-1}\|_{\mathcal{B}(\ell^2(\Gamma))}^{-2} &= \|\mathbf{F}^{-1}\|_{\mathcal{M}_n(L^\infty(\widehat{N}))}^{-2} = \left[\text{ess sup}_{\xi \in \widehat{N}} \|\mathbf{F}^{-1}(\xi)\|_2^2 \right]^{-1} \\ &= \left[\text{ess sup}_{\xi \in \widehat{N}} \lambda_{\min}^{-1} \mathbf{F}^*(\xi) \mathbf{F}(\xi) \right]^{-1} = \text{ess inf}_{\xi \in \widehat{N}} \lambda_{\min} \mathbf{F}^*(\xi) \mathbf{F}(\xi) = A_f. \end{aligned}$$

Hence $A_f > 0$, and then, having in mind (8), we prove condition (b).

Assume now (b). Since the entries of $\mathbf{F}(\xi)$ belong to $L^\infty(\widehat{N})$ and $\text{ess inf}_{\xi \in \widehat{N}} |\det \mathbf{F}(\xi)| > 0$, there exists $[\mathbf{F}^*(\xi)]^{-1}$, a.e. $\xi \in \widehat{N}$; besides, $[\mathbf{F}^*(\xi)]^{-1} \in \mathcal{M}_\kappa(L^\infty(\widehat{N}))$. Since $\mathcal{S}(L^*(\Gamma))$ is a C^* -subalgebra of $\mathcal{M}_\kappa(L^\infty(\widehat{N}))$ and \mathbf{F}^* belongs to $\mathcal{S}(L^*(\Gamma))$, its inverse $(\mathbf{F}^*)^{-1}$ also belongs to $\mathcal{S}(L^*(\Gamma))$. Hence there exists a unique $\mathbf{g} \in L^*(\Gamma)$ such that $\Lambda_{\mathbf{g}} = (\mathbf{F}^*)^{-1}$. By means of the C^* -isomorphism $\mathcal{S}\Lambda^{-1}$, we deduce that $\Lambda_{\mathbf{g}}\Lambda_{f^*} = Id$, and then

$$\sum_{\gamma \in \Gamma} \langle \mathbf{a}, L_\gamma \mathbf{f} \rangle L_\gamma \mathbf{g} = \Lambda_{\mathbf{g}}(\Lambda_{f^*} \mathbf{a}) = \mathbf{a}, \quad \mathbf{a} \in \ell^2(\Gamma).$$

From Theorem 5, the systems $\{L_\gamma \mathbf{f}\}_{\gamma \in \Gamma}$ and $\{L_\gamma \mathbf{g}\}_{\gamma \in \Gamma}$ are Bessel sequences. Hence, from [8, Theorem 3.6.6], the system $\{L_\gamma \mathbf{f}\}_{\gamma \in \Gamma}$ is a Riesz basis for $\ell^2(\Gamma)$ with dual Riesz basis $\{L_\gamma \mathbf{g}\}_{\gamma \in \Gamma}$.

Finally, since $\mathbf{F}^*(\xi)\mathbf{G}(\xi) = \mathbf{F}^*(\xi)[\mathbf{F}^*(\xi)]^{-1} = \mathbf{I}_\kappa$, a.e. $\xi \in \widehat{N}$, having in mind that $\mathcal{T}_\Gamma \mathbf{g}(\xi)$ is the first column of the matrix $\mathbf{G}(\xi)$ and Lemma 3, we deduce that \mathbf{g} is the unique element in $\ell^2(\Gamma)$ satisfying (7). \square

Acknowledgments: This work has been supported by the grant MTM2017-84098-P from the Spanish *Ministerio de Economía y Competitividad (MINECO)*.

References

- [1] A. Aldroubi, Q. Sun and W. S. Tang. Convolution, average sampling, and a Calderon resolution of the identity for shift-invariant spaces. *J. Fourier Anal. Appl.*, 11(2):215–244, 2005.
- [2] D. Barbieri, E. Hernández and J. Parcet. Riesz and frame systems generated by unitary actions of discrete groups. *Appl. Comput. Harmon. Anal.*, 39(3): 369–399, 2015.
- [3] D. Barbieri, E. Hernández and V. Paternostro. The Zak transform and the structure of spaces invariant by the action of an LCA group. *J. Funct. Anal.*, 269:1327–1358, 2015.
- [4] J. J. Benedetto. *Harmonic Analysis and Applications*. CRC Press, Boca Raton FL, 1996.
- [5] M. Benzi and P. Boito. Decay properties for functions of matrices over C^* -algebras. *Linear Algebra Appl.*, 456:174–198, 2014.
- [6] B. Blackadar. *Operator Algebras: Theory of C^* -Algebras and von Neumann Algebras*. Encyclopaedia Math. Sci., Vol. 122, Springer, Berlin, 2006.
- [7] C. de Boor, R. A. DeVore and A. Ron. On the construction of multivariate pre-wavelets. *Constr. Approx.*, 9:123–166, 1993.
- [8] O. Christensen. *An Introduction to Frames and Riesz Bases*. Second Edition. Birkhäuser, Boston, 2016.

- [9] H. R. Fernández-Morales, A. G. García, M. A. Hernández-Medina and M. J. Muñoz-Bouzo. Generalized sampling: from shift-invariant to U -invariant spaces. *Anal. Appl.*, 13(3):303–329, 2015.
- [10] G. B. Folland. *A Course in Abstract Harmonic Analysis*. CRC Press, Boca Raton FL, 1995.
- [11] A. G. García and G. Pérez-Villalón. Multivariate generalized sampling in shift-invariant spaces and its approximation properties. *J. Math. Anal. Appl.*, 355:397–413, 2009.
- [12] E. Hernández, H. Sikic, G. Weiss and E. Wilson. Cyclic subspaces for unitary representations of LCA groups; generalized Zak transform. *Colloq. Math.*, 118(1):313–332, 2010.
- [13] H. Heuser *Functional Analysis*. John Wiley, New York, 1982.
- [14] R. A. Horn and C. R. Johnson. *Matrix Analysis*. Cambridge University Press, 1999.
- [15] R. Q. Jia and C. A. Micchelli. Using the refinement equations for the construction of pre-wavelets II: Powers of two. In *Curves and Surfaces*. P. J. Laurent, Le Méhauté, L. L. Schumaker, Eds. Academic Press, Boston, pp. 209–246, 1991.
- [16] J. M. Kim and K. H. Kwon. Sampling expansion in shift invariant spaces. *Int. J. Wavelets Multiresolut. Inf. Process.*, 6(2):223–248, 2008.
- [17] J. J. Lei, R. Q. Jia, and E. W. Cheney. Approximation from shift-invariant spaces by integral operators. *SIAM J. Math. Anal.*, 28(2):481–498, 1997.
- [18] A. L. González and M. C. Moure.. Crystallographic Haar wavelets. *J. Fourier Anal. Appl.*, 17(6), 1119–1137, 2011.
- [19] K. Guo, D. Labate, W. Q. Lim, G. Weiss and E. Wilson. The theory of wavelets with composite dilations. In *Harmonic Analysis and Applications*, C. Heil (Ed.), pp. 231–250, Birkhäuser, Boston, 2006.
- [20] J. MacArthur and K. Taylor. Wavelets with crystal symmetry shifts. *J. Fourier Anal. Appl.*, 17(6), 1109–1118, 2011.
- [21] B. Manning. Composite multi resolution analysis wavelets. PhD Thesis, <http://openscholarship.wustl.edu/etd/716/>, 2012.
- [22] M. A. Rieffel. Actions of finite groups on C^* -algebras. *Math. Scand.*, 47(1), 157–176, 1980.
- [23] A.A. Selvan and R. Radha. Sampling and reconstruction in shift-invariant spaces on \mathbb{R}^d . *Ann. Mat. Pura Appl.*, 194(6), 1683–1706, 2015.
- [24] M. Takesaki. *Theory of Operator Algebras I*. Encyclopaedia Math. Sci., Vol. 124, Springer, Berlin, 2002.
- [25] K.F. Taylor. C^* -algebras of crystal groups. *Operator Theory: Advances and Applications*, 41: 511–518, 1989.

- [26] P. P. Vaidyanathan. *Multirate Systems and Filter Banks*. Prentice Hall, 1993.
- [27] X. Zhou and W. Sun. On the sampling theorem for wavelet subspaces. *J. Fourier Anal. Appl.*, 5(4):347–354, 1999.