

CANONICAL FORMS FOR CONGRUENCE OF MATRICES: A TRIBUTE TO H. W. TURNBULL AND A. C. AITKEN*

FERNANDO DE TERÁN†

Abstract. A canonical form for congruence of matrices was introduced by Turnbull and Aitken in 1932. More than 70 years later, in 2006, another canonical form for congruence has been introduced by Horn and Sergeichuk. The main purpose of this paper is to compare both canonical forms and provide a brief survey on the history of the canonical form for congruence.

Key words. matrices, canonical forms, congruence, matrix pencils

AMS subject classifications. 15A21, 15A22

1. Introduction.

1.1. Congruence and similarity. We consider the following two actions of the general linear group of $n \times n$ invertible matrices with complex entries (denoted by $\mathrm{GL}(n, \mathbb{C})$) on the set of $n \times n$ matrices with complex entries (denoted by $\mathbb{C}^{n \times n}$):

$$\begin{array}{ccc} \mathrm{GL}(n, \mathbb{C}) \times \mathbb{C}^{n \times n} & \longrightarrow & \mathbb{C}^{n \times n} \\ (P, A) & \longmapsto & PAP^{-1} \end{array}, \quad \text{and} \quad \begin{array}{ccc} \mathrm{GL}(n, \mathbb{C}) \times \mathbb{C}^{n \times n} & \longrightarrow & \mathbb{C}^{n \times n} \\ (P, A) & \longmapsto & PAP^T. \end{array}$$

The first one is known as the action of *similarity* and the second one as the action of *congruence*. Accordingly, two matrices $A, B \in \mathbb{C}^{n \times n}$ are said to be *similar* if there is a nonsingular $P \in \mathbb{C}^{n \times n}$ such that $PAP^{-1} = B$ and they are said to be *congruent* if there is a nonsingular P such that $PAP^T = B$. Similar matrices correspond to the same linear transformation in different bases, whereas congruent matrices correspond to equivalent bilinear forms. Both the similarity and the congruency determine equivalence relations in $\mathbb{C}^{n \times n}$. These relations allow us to classify the set $\mathbb{C}^{n \times n}$ into equivalence classes (also called *orbits*) consisting of equivalent matrices. A usual approach to identify these equivalence classes is by introducing a canonical form for the corresponding relation. Then the problem of determining whether or not two matrices are equivalent reduces to the problem of finding the canonical form of these matrices. There are several interesting connections between canonical forms for similarity and congruence. For instance, the *Sylvester equation* $XA - AX = 0$ can be solved in terms of the canonical form for similarity of A [5, Chapter VIII, §2], and the same holds for the matrix equation $XA + AX^T = 0$ with the canonical form for congruence [4]. But there is a remarkable difference between them. On the one hand, the canonical form for similarity goes back to Jordan (1838-1922) and it is widely known by the mathematical community (it is the famous *Jordan Canonical Form*, from now on, the JCF). Moreover, it is completely accepted as a canonical form par excellence. For instance, it is studied by first-course degree students in mathematics and in other scientific disciplines, and most textbooks in basic (and not so basic) linear algebra contain it. On the contrary, the canonical form for congruence was introduced in 1932 and it is not sufficiently known even by linear algebra researchers. Moreover, several canonical forms for congruence have been proposed later by different authors and none of them

*Partially supported by the Ministerio de Educación y Ciencia of Spain through grant MTM-2009-09281

†Departamento de Matemáticas, Universidad Carlos III de Madrid, Avda. Universidad 30, 28911 Leganés, Spain fteran@math.uc3m.es

is still completely accepted by researchers (this would not be surprising, since two of them have been introduced in 2006). The canonical form for congruence is the subject of the present paper.

1.2. Some history about the canonical form for congruence. A canonical form for congruence of complex matrices was derived by Turnbull and Aitken in a classical reference on matrix theory [18, p. 139]. Turnbull and Aitken obtained first a canonical form for congruence of pencils of the type $\mu A + \lambda A^T$ (here μ, λ are complex variables) and, from this canonical form they derive the canonical form for matrices just by taking $\mu = 1, \lambda = 0$. The most immediate difference between the JCF and the congruence canonical form of Turnbull and Aitken (from now on, the *TA canonical form*) is that whereas the JCF consists of just one type of blocks (the *Jordan blocks* associated with a particular eigenvalue) the *TA canonical form* consists of six types of blocks (see Section 3).

In 1947, Hodge and Pedoe, without mentioning the work of Turnbull and Aitken at all, derived in [7] the same canonical form as Turnbull and Aitken (except for a slight difference in the presentation of one type of blocks). Their canonical form for $A \in \mathbb{C}^{n \times n}$ comes from the canonical reduction of matrix pairs (R, S) , with R symmetric and S skew-symmetric, applied to the symmetric and skew-symmetric parts of A . This reduction is obtained by the authors using similar techniques as the ones introduced by Bromwich forty years before in [1] for pairs of symmetric matrices.

A long time after the publication of Turnbull and Aitken's book (we are using a reprinted edition [18] of the original book, which goes back to 1932), Sergeichuk provided in [16] canonical forms for congruence over any field \mathbb{F} of characteristic not 2 up to classification of Hermitian forms over finite extensions of \mathbb{F} . Based on that work Horn and Sergeichuk established in [8] a canonical form for congruence over fields of characteristic not two with an involution (later extended in [11] to bilinear forms over arbitrary algebraically closed fields and real closed fields). When particularized to the complex field, they achieve in [10] another canonical form for congruence in $\mathbb{C}^{n \times n}$. From now on, we will refer to it as the *HS canonical form*. The types of blocks in this canonical form are different than the types of blocks in the *TA canonical form*. In particular, the *HS canonical form* consists of three types of blocks instead of the six types in the *TA form*. This lower number of blocks makes this form more manageable and it is, for instance, the form that has been used to solve the equation $XA + AX^T = 0$ in [4]. In the meantime, another canonical form was proposed in [3]. Nonetheless, as it was noticed by Horn and Sergeichuk in [10, p. 1013], some of the canonical matrices in this form are "cumbersome and not canonical".

In 1992 Thompson published an expository paper [17], where he showed canonical forms for congruence of pencils $\rho R + \sigma S$, with R, S being symmetric or skew-symmetric (both real and complex). The case where R is symmetric and S skew-symmetric is closely related to the type $\mu A + \lambda A^T$. More precisely, as noticed by Turnbull and Aitken [18, p. 136], an elementary change of variables leads the pencil $\rho R + \sigma S$ to the pencil $\mu A + \lambda A^T$. In 1996, apparently unaware of [18] and [7], Lee and Weinberg [13] followed Thompson's paper to derive a canonical form for congruence of matrices from the canonical form of pencils $\rho R + \sigma S$, with R symmetric and S skew-symmetric. Their approach is the same as the one of Hodge and Pedoe. These authors focus only on the real case, and they just mention that a similar procedure can be followed for complex matrices. This procedure should lead to the *TA canonical form*. This is precisely the approach followed by Schröder [15] and he actually obtains six types of blocks. Nonetheless, these blocks have a different appearance than the ones of the

TA canonical form due to some further modifications addressed by the author. This amounts to, at least, three canonical forms for congruence after Turnbull and Aitken's one (without considering the one of Lee and Weienberg).

The most surprising fact in all this story is that none of the papers [13, 16, 3, 8, 10, 15] mention the work of Turnbull and Aitken. This is actually more striking because most of them cite Thompson's paper [17], where Turnbull and Aitken's book is included among the references (though it is just cited as a reference for historical remarks [17, pp. 335, 358]). One of the main purposes of the present paper is to point out the work of Turnbull and Aitken. On the other hand, we think that the HS canonical form is the simplest one among all canonical forms mentioned above (including the TA canonical form). Hence, we consider interesting to establish the relationship between this form and the pioneer one of Turnbull and Aitken. Since both forms provide reductions under congruence for matrices in $\mathbb{C}^{n \times n}$, there should be a correspondence between blocks in each of these canonical forms. To explicitly state this correspondence is the other main goal of this paper.

One of the advantages of the TA canonical form of A is that it immediately gives the spectral structure of the associated pencil $\mu A + \lambda A^T$. These pencils are known nowadays as *T-palindromic* pencils, though this name has been introduced recently¹. They are becoming a subject of interest in the linear algebra community mostly motivated by its appearance in several applied problems [6, 14, 12]. As noted before, the *T-palindromic* pencils $\mu A + \lambda A^T$ are in one-to-one correspondence with pencils of the form $\rho R + \sigma S$, with R symmetric and S skew-symmetric. Horn and Sergeichuk, proceeding in the opposite way as Turnbull and Aitken, derive canonical forms for congruence of these last kind of pencils from their canonical form for congruence of matrices. Nonetheless, the canonical form of Horn and Sergeichuk does not explicitly show the spectral structure of the pencil. In the present paper, we will show how the spectral structure of the associated palindromic pencil $\mu A + \lambda A^T$ can be recovered from the HS canonical form of A .

We close this Introduction by noticing that this is not the first time that the book of Turnbull and Aitken is "rediscovered". Brualdi published in [2] an "old" proof for the JCF whose basic idea was found in that book. This leads us to ask : What other pearls may be hidden in this old book?

2. Spectral structure of *T-palindromic* matrix pencils. We recall in this section some basic notions of matrix pencils that will be used in the proof of our main result (Theorem 3.5). Only in this section we allow for rectangular matrices $A, B \in \mathbb{C}^{m \times n}$. We will say that two matrix pencils $\mu A + \lambda B$ and $\mu C + \lambda D$ are *equivalent* if there exist two nonsingular matrices $P \in \mathbb{C}^{m \times m}$ and $Q \in \mathbb{C}^{n \times n}$ such that

$$P(\mu A + \lambda B)Q = \mu C + \lambda D.$$

Each matrix pencil $\mu A + \lambda B$ is equivalent to a block diagonal pencil of the form

$$\text{diag}(J, N, L_{\varepsilon_1}, \dots, L_{\varepsilon_p}, L_{\eta_1}^T, \dots, L_{\eta_q}^T), \quad (2.1)$$

where

$$J = \text{diag}(\mu J_{n_1}(\lambda_1) + \lambda I_{n_1}, \dots, \mu J_{n_r}(\lambda_r) + \lambda I_{n_r}), \quad (2.2)$$

¹It was introduced privately by D. Steven Mackey in 2003, but the first reference where this name appears is [6]

and

$$N = \text{diag}(\mu I_{m_1} + \lambda J_{m_1}(0), \dots, \mu I_{m_s} + \lambda J_{m_s}(0)), \quad (2.3)$$

with

$$J_d(\lambda_i) = \begin{bmatrix} \lambda_i & 1 & & & \\ & \ddots & \ddots & & \\ & & \ddots & \ddots & \\ & & & \lambda_i & 1 \\ & & & & \lambda_i \end{bmatrix}$$

being a $d \times d$ Jordan block associated with $\lambda_i \in \mathbb{C}$. Also

$$L_{\varepsilon_i} = \begin{bmatrix} \mu & \lambda & & & \\ & \ddots & \ddots & & \\ & & \ddots & \ddots & \\ & & & \mu & \lambda \end{bmatrix}_{\varepsilon_i \times (\varepsilon_i + 1)}$$

is an $\varepsilon_i \times (\varepsilon_i + 1)$ right singular block, and $L_{\eta_i}^T$ is an $(\eta_i + 1) \times \eta_i$ left singular block. The pencils (2.2) and (2.3) comprise the *regular structure* (finite and infinite, respectively) of $\mu A + \lambda B$, and the remaining diagonal blocks of (2.1) comprise the so-called *singular structure*.

The numbers $-\lambda_1, \dots, -\lambda_r$ are the *finite eigenvalues* of $\mu A + \lambda B$, and it is said that $\mu A + \lambda B$ has an *infinite eigenvalue* if N in (2.1) is nonempty. Accordingly, we will refer to the diagonal blocks of N as *Jordan blocks associated with the infinite eigenvalue*.

The numbers η_1, \dots, η_q and $\varepsilon_1, \dots, \varepsilon_p$ are known as, respectively, the *left and right minimal indices* of $\mu A + \lambda B$. When $\mu A + \lambda B$ is square (and, in particular, when it is T -palindromic), we have $p = q$.

The pencil (2.1) is known as the *Kronecker Canonical Form* (KCF) of $\mu A + \lambda B$, and it is unique up to permutation of the diagonal blocks. This canonical form comprises the so-called *spectral structure* of $\mu A + \lambda B$, consisting of the eigenvalues of $\mu A + \lambda B$, together with the sizes of the associated Jordan blocks, and the minimal indices. For a proof of the existence and uniqueness of the KCF we refer the reader to the classical reference on matrix theory [5, Chapter XII], though pencils in that book are written as $A + \lambda B$ (that is, with just one variable) instead of $\mu A + \lambda B$.

When particularizing to the case of T -palindromic matrix pencils the spectral structure has several restrictions imposed by the particular structure of this kind of pencils. These constraints are fully explained in [17], but they are present in the book by Turnbull and Aitken. Following these two references we state the spectral restrictions in the theorem below.

THEOREM 2.1. *Let $\mu A + \lambda A^T$ be a T -palindromic pencil. Then, the KCF of $\mu A + \lambda A^T$ is subject to the following restrictions.*

- (i) *The left and right minimal indices of $\mu A + \lambda A^T$ coincide, that is: if ε is a right minimal index of $\mu A + \lambda A^T$ then it is also a left minimal index and vice versa.*
- (ii) *Each Jordan block with odd size associated with the eigenvalue $\lambda = 1$ occurs an even number of times.*
- (iii) *Each Jordan block with even size associated with the eigenvalue $\lambda = -1$ occurs an even number of times.*
- (iv) *The Jordan blocks associated with eigenvalues $\lambda \neq \pm 1$ occur in pairs: $J_s(\lambda), J_s(1/\lambda)$ (here we understand that the eigenvalue 0 is paired up with the infinite eigenvalue).*

3. The canonical structure for congruence. In this section we will follow the notation from [11]

$$[A \setminus B] := \begin{bmatrix} 0 & B \\ A & 0 \end{bmatrix}.$$

We begin by introducing the canonical form of Turnbull and Aitken.

THEOREM 3.1. (TA canonical form of congruence) [18, p. 139] *Each matrix $A \in \mathbb{C}^{n \times n}$ is congruent to a direct sum of blocks of the following types:*

$$\begin{aligned} \text{(a) } \mathcal{A}_{2k+1} &= \left[\begin{bmatrix} 0 & 1 \\ & \ddots & \ddots \\ & & 0 & 1 \end{bmatrix} \setminus \begin{bmatrix} 1 & & & \\ & \ddots & & \\ & & \ddots & 1 \\ & & & 0 \end{bmatrix} \right]_{(2k+1) \times (2k+1)} \\ \text{(b) } \mathcal{B}_{2k}(c) &= \left[\begin{bmatrix} 0 & & c \\ & c & 1 \\ & \ddots & \ddots \\ c & 1 & 0 \end{bmatrix} \setminus \begin{bmatrix} & & & 1 \\ & & & c \\ & & 1 & \\ & & \ddots & \\ 1 & c & & 0 \end{bmatrix} \right]_{2k \times 2k}, \quad c \neq \pm 1. \\ \text{(c) } \mathcal{C}_{2k+1} &= \begin{bmatrix} 0 & & & & & & & & & 1 \\ & & & & & & & & & 1 \\ & & & & & & & & & \ddots \\ & & & & & & & & & \ddots \\ & & & & & & & & & 1 \\ & & & & & & & & & 1 \\ & & & & & & & & & \ddots \\ & & & & & & & & & \ddots \\ & & & & & & & & & 1 \\ & & & & & & & & & 1 \\ & & & & & & & & & \ddots \\ & & & & & & & & & \ddots \\ & & & & & & & & & 1 \\ & & & & & & & & & 1 \\ & & & & & & & & & \ddots \\ & & & & & & & & & \ddots \\ & & & & & & & & & 0 \end{bmatrix}_{(2k+1) \times (2k+1)} \\ \text{(d) } \mathcal{D}_{2k} &= \left[\begin{bmatrix} 0 & & & 1 \\ & & & -1 \\ & & & \ddots \\ & & & 0 \\ 1 & -1 & \ddots & \end{bmatrix} \setminus \begin{bmatrix} & & & 1 \\ & & & 1 \\ & & 1 & \\ & & \ddots & \\ 1 & 1 & & 0 \end{bmatrix} \right]_{2k \times 2k} \quad (k \text{ even}). \\ \text{(e) } \mathcal{E}_{2k} &= \begin{bmatrix} 0 & & & & & & & & & 1 \\ & & & & & & & & & 1 \\ & & & & & & & & & \ddots \\ & & & & & & & & & \ddots \\ & & & & & & & & & 1 \\ & & & & & & & & & 1 \\ & & & & & & & & & \ddots \\ & & & & & & & & & \ddots \\ & & & & & & & & & 1 \\ & & & & & & & & & 1 \\ & & & & & & & & & \ddots \\ & & & & & & & & & \ddots \\ & & & & & & & & & 0 \end{bmatrix}_{2k \times 2k} \\ \text{(f) } \mathcal{F}_{2k} &= \left[\begin{bmatrix} 0 & & & -1 \\ & & & 1 \\ & & & \ddots \\ & & & 0 \\ -1 & 1 & \ddots & \end{bmatrix} \setminus \begin{bmatrix} 0 & & & 1 \\ & & & 1 \\ & & 1 & \\ & & \ddots & \\ 1 & 1 & & 0 \end{bmatrix} \right]_{2k \times 2k} \quad (k \text{ odd}). \end{aligned}$$

Now we will state the HS canonical form. For this we introduce the $k \times k$ matrix

$$\Gamma_k = \begin{bmatrix} 0 & & & & & & & & & (-1)^{k+1} \\ & & & & & & & & & (-1)^k \\ & & & & & & & & & \ddots \\ & & & & & & & & & -1 \\ & & & & & & & & & \ddots \\ & & & & & & & & & 1 \\ & & & & & & & & & 1 \\ & & & & & & & & & -1 \\ & & & & & & & & & -1 \\ & & & & & & & & & 1 \\ & & & & & & & & & 0 \end{bmatrix} \quad (\Gamma_1 = [1])$$

and, for each $c \in \mathbb{C}$, the $2k \times 2k$ matrix

$$H_{2k}(c) = [J_k(c) \setminus I_k] \quad (H_2(c) = [c \setminus 1]),$$

where $J_k(c)$ is a $k \times k$ Jordan block associated with c .

The HS canonical form is given in the following theorem.

THEOREM 3.2. (HS canonical form for congruence) [10, Theorem 1.1] *Each square complex matrix is congruent to a direct sum, uniquely determined up to permutation of summands, of canonical matrices of the three types*

Type 0	$J_k(0)$	(3.1)
Type I	Γ_k	
Type II	$H_{2k}(c)$, $0 \neq c \neq (-1)^{k+1}$ <i>c is determined up to replacement by c^{-1}</i>	

Comparing the statements of these theorems one immediately realizes on a striking difference. Unlike the HS canonical form, the TA canonical form is not claimed to be unique. The difference in the number of canonical blocks in these two canonical forms may lead to think that there are some redundant blocks in the TA canonical form. But, by comparing blocks in the two canonical forms, we will conclude that this canonical form is indeed unique (up to permutation of blocks), and that there are no redundant blocks.

In order to prove our main result, we will show first how to recover the spectral structure of the pencil $\mu A + \lambda A^T$ from the HS canonical form of A .

THEOREM 3.3. *The KCF of the T -palindromic pencil $\mu A + \lambda A^T$ can be recovered from the HS canonical form of A as follows:*

- (i) $\mu J_{2k}(0) + \lambda J_{2k}(0)^T$ is equivalent to $(\mu J_k(0) + \lambda I_k) \oplus (\mu I_k + \lambda J_k(0))$.
- (ii) $\mu J_{2k+1}(0) + \lambda J_{2k+1}(0)^T$ is equivalent to $L_{2k} \oplus L_{2k}^T$.
- (iii) $\mu H_{2k}(c) + \lambda H_{2k}(c)^T$ is equivalent to $(\mu J_k(c) + \lambda I_k) \oplus (\mu J_k(1/c) + \lambda I_k)$ ($c \neq 0$).
- (iv) $\mu \Gamma_k + \lambda \Gamma_k^T$ is equivalent to $\mu J_k((-1)^{k+1}) + \lambda I_k$.

Proof. Since $J_{2k}(0)$ is congruent to $H_{2k}(0)$ [16, p. 493], we conclude that $\mu J_{2k}(0) + \lambda J_{2k}(0)^T$ is equivalent to $\mu H_{2k}(0) + \lambda H_{2k}(0)^T$ and, by the uniqueness of the KCF, this is in turn equivalent to $(\mu J_k(0) + \lambda I_k) \oplus (\mu I_k + \lambda J_k(0))$. This proves (i).

Since $J_{2k+1}(0)$ is congruent to \mathcal{A}_{2k+1} [16, p. 492] claim (ii) follows easily.

Claim (iii) is immediate from the uniqueness of the KCF. Finally, to prove (iv) we multiply

$$\Gamma_k^{-T} (\mu \Gamma_k + \lambda \Gamma_k^T) = \mu \begin{bmatrix} (-1)^{k+1} & 2 & & \star \\ & \ddots & \ddots & \\ & & (-1)^{k+1} & 2 \\ 0 & & & (-1)^{k+1} \end{bmatrix} + \lambda I_k$$

where the symbol \star denotes entries with no relevance to the argument. Now (iv) follows again by the uniqueness of the KCF. \square

Notice that the KCF of $\mu A + \lambda A^T$ obtained in Theorem 3.3 is coherent with the restrictions stated in Theorem 2.1.

Since congruence of matrix pencils is a particular case of equivalence, congruent matrix pencils have the same KCF. Turnbull and Aitken obtained in [18, pp. 138-139] a canonical form for congruence of T -palindromic matrix pencils that exhibits the KCF of the pencil $\mu A + \lambda A^T$. From this canonical form they derive the canonical

form for congruence of matrices just by taking $\mu = 1, \lambda = 0$. As a consequence, the TA canonical form of A allows to know the KCF of $\mu A + \lambda A^T$. On the other hand, Theorem 3.3 shows also how to recover the KCF of $\mu A + \lambda A^T$ from the HS canonical form. If we were able to prove that, conversely, the KCF of $\mu A + \lambda A^T$ uniquely determines the congruence class of A then we could establish the relationship between the HS and the TA canonical forms. The following Lemma shows that this is indeed possible.

LEMMA 3.4. *Two square matrices $A, B \in \mathbb{C}^{n \times n}$ are congruent if and only if the pencils $\mu A + \lambda A^T$ and $\mu B + \lambda B^T$ are equivalent.*

Proof. If A, B are congruent then the pencils $\mu A + \lambda A^T$ and $\mu B + \lambda B^T$ are clearly equivalent. Let us prove the converse. Assume that $\mu A + \lambda A^T$ and $\mu B + \lambda B^T$ are equivalent. Then, there exists two nonsingular matrices P, Q such that $PAQ = B$ and $PA^TQ = B^T$, so

$$P \left(\frac{A + A^T}{2} \right) Q = \frac{B + B^T}{2} \quad \text{and} \quad P \left(\frac{A - A^T}{2} \right) Q = \frac{B - B^T}{2}.$$

Now, by Lemma III in [18, p. 130], there is a nonsingular $H \in \mathbb{C}^{n \times n}$ such that

$$H^T \left(\frac{A + A^T}{2} \right) H = \frac{B + B^T}{2} \quad \text{and} \quad H^T \left(\frac{A - A^T}{2} \right) H = \frac{B - B^T}{2},$$

and the result follows. \square

Now we are in the position to state the main result of this paper.

THEOREM 3.5. *Each of the blocks in the TA canonical form is congruent to one of the blocks in the HS canonical form as follows.*

- 1) \mathcal{A}_{2k+1} is congruent to $J_{2k+1}(0)$.
- 2) $\mathcal{B}_{2k}(c)$ is congruent to
 - 2a) $J_{2k}(0)$, if $c = 0$, and
 - 2b) $H_{2k}(c)$, if $c \neq 0$.
- 3) \mathcal{C}_{2k+1} is congruent to Γ_{2k+1} .
- 4) \mathcal{D}_{2k} is congruent to $H_{2k}(1)$ (k even).
- 5) \mathcal{E}_{2k} is congruent to Γ_{2k} .
- 6) \mathcal{F}_{2k} is congruent to $H_{2k}(-1)$ (k odd).

Proof. The result is an immediate consequence of Theorem 3.3, Lemma 3.4, and the relationship between canonical blocks in the TA canonical form of A and the KCF of $\mu A + \lambda A^T$, given by [18, pp. 135–139]

- $\mu \mathcal{A}_{2k+1} + \lambda \mathcal{A}_{2k+1}^T$ is equivalent to $L_{2k} \oplus L_{2k}^T$.
- $\mu \mathcal{B}_{2k+1}(c) + \lambda \mathcal{B}_{2k+1}(c)^T$ is equivalent to $(\mu J_k(c) + \lambda I_k) \oplus (\mu J_k(1/c) + \lambda I_k)$ ($0 \neq c \neq \pm 1$).
- $\mu \mathcal{B}_{2k+1}(0) + \lambda \mathcal{B}_{2k+1}(0)^T$ is equivalent to $(\mu J_k(0) + \lambda I_k) \oplus (\mu I_k + \lambda J_k(0))$.
- $\mu \mathcal{C}_{2k+1} + \lambda \mathcal{C}_{2k+1}^T$ is equivalent to $(\mu J_{2k+1}(1) + \lambda I_{2k+1})$.
- $\mu \mathcal{D}_{2k+1} + \lambda \mathcal{D}_{2k+1}^T$ is equivalent to $(\mu J_k(1) + \lambda I_k) \oplus (\mu J_k(1) + \lambda I_k)$.
- $\mu \mathcal{E}_{2k} + \lambda \mathcal{E}_{2k}^T$ is equivalent to $(\mu J_{2k}(-1) + \lambda I_{2k})$.
- $\mu \mathcal{F}_{2k} + \lambda \mathcal{F}_{2k}^T$ is equivalent to $(\mu J_k(-1) + \lambda I_k) \oplus (\mu J_k(-1) + \lambda I_k)$.

\square

Theorem 3.5 also allows us to explain the difference in the number of blocks between the TA and the HS canonical forms. This difference comes from the size of type 0 and type I blocks and also on the values associated with type II blocks in the HS canonical form. More precisely, type 0 blocks give rise to two different types of

Despite this redundancy in the canonical form for T -palindromic pencils, when the authors derive the canonical form for matrices by taking $\mu = 1$, $\lambda = 0$, the redundant blocks disappear without any further comment.

Acknowledgments. The author wishes to thank Froilán M. Dopico for posing the main question addressed in the paper, namely, to state the relationship between the TA and the HS canonical forms for congruence.

REFERENCES

- [1] T. J. BROMWICH, *Quadratic forms and their classification by means of invariant factors*, Cambridge University Press, Cambridge, 1906.
- [2] R. A. BRUALDI, *The Jordan Canonical Form: an Old Proof*, Amer. Math. Monthly, 94 (1987) 257–267.
- [3] B. CORBAS, G. D. WILLIAMS, *Bilinear forms over an algebraically closed field*, J. Pure Appl. Algebra 165 (3) (2001) 225–266.
- [4] F. DE TERÁN, F. M. DOPICO, *The solution of the equations $XA + AX^T = 0$ and its application to the theory of orbits*, in preparation.
- [5] F. R. GANTMACHER, *The theory of matrices*, Chelsea, New York, 1959.
- [6] A. HILLIGES, C. MEHL, AND V. MEHRMANN, *On the solution of palindromic eigenvalue problems*, in Proceedings of the 4th European Congress on Computational Methods in Applied Sciences and Engineering (ECCOMAS), JyvÅaskylÅa, Finland, 2004. CD-ROM.
- [7] W. V. D HODGE, D. PEDOE, *Methods of Algebraic Geometry*, Cambridge University Press, Cambridge, 1947.
- [8] R. A. HORN, V. V. SERGEICHUK, *Congruence of a square matrix and its transpose*, Linear Algebra Appl., 389 (2004) 347–353.
- [9] R. A. HORN, V. V. SERGEICHUK, *A regularization algorithm for matrices of bilinear and sesquilinear forms*, Linear Algebra Appl., 412 (2006) 380–395.
- [10] R. A. HORN, V. V. SERGEICHUK, *Canonical forms for complex matrix congruence and *congruence*, Linear Algebra Appl., 416 (2006) 1010–1032.
- [11] R. A. HORN, V. V. SERGEICHUK, *Canonical matrices of bilinear and sesquilinear forms*, Linear Algebra Appl., 428 (2008) 193–223.
- [12] T.-M. HUANG, W.-W. LIN, AND J. QIANG, *Structure-preserving algorithms for palindromic quadratic eigenvalue problems arising from vibration of fast trains*, SIAM J. Matrix Anal. Appl., 30 (4) (2009) 1566–1592.
- [13] J. M. LEE, D. A. WEINBERG, *A note on the canonical form for matrix congruence*, Linear Algebra Appl., 249 (1996) 207–215.
- [14] D. S. MACKEY, N. MACKEY, C. MEHL, AND V. MEHRMANN, *Structured polynomial eigenvalue problems: Good vibrations from good linearizations*, SIAM J. Matrix Anal. Appl., 28 (2006) 1029–1051.
- [15] C. SCHRÖDER, *A canonical form for palindromic pencils and palindromic factorizations*, preprint. Available at http://www.matheon.de/preprints/3822_Sch06.pdf
- [16] V. V. SERGEICHUK, *Classification problems for system of forms and linear mappings*, Math. USSR, Izvestiya 31 (3) (1998) 481–501.
- [17] R. C. THOMPSON, *Pencils of Complex and Real Symmetric Matrices*, Linear Algebra Appl., 147 (1991) 323–371.
- [18] H. W. TURNBULL, A. C. AITKEN, *An introduction to the Theory of Canonical Matrices*, Dover, New York, 1961 (First published by: Blackie & Son Limited, 1932).