

A GEOMETRIC DESCRIPTION OF THE SETS OF PALINDROMIC AND ALTERNATING MATRIX PENCILS WITH BOUNDED RANK*

FERNANDO DE TERÁN[†]

Abstract. The sets of $n \times n$ \mathbb{T} -palindromic, \mathbb{T} -anti-palindromic, \mathbb{T} -even, and \mathbb{T} -odd matrix pencils with rank at most $r < n$ are algebraic subsets of the set of $n \times n$ matrix pencils. In this paper, we determine their dimension and we prove that they are all irreducible. This is in contrast with the non-structured case, since it is known that the set of $n \times n$ matrix pencils with rank at most $r < n$ is an algebraic set with $r + 1$ irreducible components. We also show that these sets of structured pencils with bounded rank are the closure of the congruence orbit of a certain structured pencil given in canonical form. This allows us to determine the generic canonical form of a structured $n \times n$ matrix pencil with rank at most r , for any of the previous structures.

Key words. Matrix pencil, \mathbb{T} -palindromic, \mathbb{T} -alternating, strict equivalence, congruence, orbit, spectral information, algebraic set

AMS subject classifications. 15A22, 15A18, 15A21, 65F15

1. Introduction. Structured matrix pencils arise in several problems of applied mathematics, either as modeling physical systems by themselves or as a tool to compute the relevant information in the analysis of other higher-order systems by linearization (see, for instance, [22] and the references therein). Some of the structures appearing most frequently in applications are the skew-symmetric, the \mathbb{T} -palindromic (or \mathbb{T} -anti-palindromic), and the \mathbb{T} -alternating structures (see Section 2 for the definition of these structures). Among the sets of $n \times n$ structured matrix pencils, an important class from the point of view of applications is the class of *low-rank pencils*. In this paper, low-rank means essentially rank-deficient, that is, we are interested in $n \times n$ pencils with rank r and $r < n$. Low-rank pencils arise when modeling systems that depend on many parameters, but only a few of them are modified (or perturbed), regardless of the size (in norm) of the modification. Some particular settings where low-rank pencils arise include dissipative dynamical systems [2, §1.2], network analysis in electrical engineering [27], or multi-body system simulation [18].

Therefore, low-rank matrix pencils naturally arise associated with low-rank perturbations, a subject which has attracted the attention of researchers in the recent years [1, 2, 4, 7, 9, 24, 25]. In a similar way as the understanding of the underlying geometry of the set of matrix pencils is helpful to analyze the change of the scalar spectral information under small perturbations [16, 17], the analysis of the geometry of low-rank matrix pencils may be helpful in the explanation of the change of the spectral information under low-rank perturbations. To be more precise, the scalar spectral information (*partial multiplicities* and *minimal indices*) comprises the invariants of matrix pencils under strict equivalence transformations. Two matrix pencils $A_0 + \lambda A_1$ and $B_0 + \lambda B_1$ are said to be *strictly equivalent* if there exist two nonsingular matrices V, W such that $V(A_0 + \lambda A_1)W = B_0 + \lambda B_1$. Then, all matrix pencils having the same scalar spectral information lie in the *strict equivalence orbit* of a given pencil

***Funding:** This work was supported by the Ministerio de Economía y Competitividad of Spain through grants MTM2015-68805-REDT, and MTM2015-65798-P, and by the Ministerio de Educación, Cultura y Deportes of Spain through grant PRX16/00128 “Programa de estancias de movilidad de profesores e investigadores en centros de enseñanza superior e investigación ”Salvador de Madariaga””

[†]Departamento de Matemáticas, Universidad Carlos III de Madrid, Avda. Universidad 30, 28911 Leganés, Spain. fteran@math.uc3m.es.

$A_0 + \lambda A_1$. As a consequence, the knowledge of the geometry of matrix pencils, in terms of the inclusion relationships between the closures of strict equivalence orbits, may allow to explain the changes in the spectral information due to small perturbations. This is translated to the set of low-rank matrix pencils and low-rank perturbations.

A description of the set of low-rank pencils (non-structured) was provided in [4], where it was proved that the set of $m \times n$ matrix pencils with rank at most $r < \min\{m, n\}$ is an algebraic set with $r + 1$ irreducible components, which are the closures of the equivalence orbits of some specific pencils given in Kronecker canonical form (a similar description has been recently obtained in [11, Th. 3.2] for the set of $m \times n$ matrix polynomials with grade d and rank at most r). Another description of the set of $m \times n$ matrix pencils with rank at most $r < \min\{m, n\}$ has been presented in the recent papers [7, 8]. In [7, Lemma 3.1], the authors provide a decomposition of the set of $n \times n$ matrix pencils with rank at most r as the union of $r + 1$ sets consisting of sums of r rank-1 pencils in such a way that some of the column vectors of this sum are constant vectors, and the remaining ones are allowed to have degree 1 (though in [7] it is only stated for square pencils, such a decomposition is valid also for rectangular $m \times n$ pencils with rank $r < \min\{m, n\}$). Then, it has been proved in [8] that each of these sets corresponds to each of the irreducible components of the algebraic set of matrix pencils with rank at most r , which results in a more constructible description of these irreducible components.

If we restrict ourselves to structured pencils and, accordingly, to structured perturbations, some remarkable differences arise. The restrictions imposed by the structure may lead to a different generic behavior under low rank perturbations [10]. This is just an indication that the analysis of the set of structured low-rank pencils deserves some special attention. In the recent years, some effort has been devoted to analyze and describe the geometry of structured matrix pencils. In particular, the set of skew-symmetric pencils has been studied with detail in [13, 14], and the set of symmetric pencils has been analyzed in [15]. However, no special attention had been paid so far to the set of structured matrix pencils with bounded rank. Up to our knowledge, the only reference on this is the recent work [12]. In that paper, the authors describe the generic scalar spectral information of skew-symmetric matrix polynomials with bounded rank using orbit closures. This includes the case of skew-symmetric matrix pencils. The main results in the present work are the counterpart of those in [12], but for \top -palindromic, \top -anti-palindromic, and \top -alternating matrix pencils, instead of skew-symmetric ones.

The set of $n \times n$ structured matrix pencils with rank at most $r < n$, denoted by \mathbb{S}_r , for any of the structures in Table 2.1, is an algebraic subset of the set of pencils $A_0 + \lambda A_1$, which can be identified with \mathbb{C}^{2n^2} (by considering a pencil $A_0 + \lambda A_1$ as a pair (A_0, A_1)). To see this, notice that \mathbb{S}_r is defined as the intersection of two algebraic sets. The first one is the set defined by the specific structure, and the second one is the set defined by the low-rank condition. Both them are algebraic sets because they can be defined in terms of multivariable polynomials in the entries of the pencil, in other words, in terms of multivariable polynomials with $2n^2$ variables (the n^2 coordinates of A_0 and the n^2 coordinates of A_1). For the second set, these polynomials are all the $(r + 1) \times (r + 1)$ minors of general $n \times n$ pencils. For the first set, the polynomials depend on the structure. For instance, for the \top -palindromic structure, they are $[A_0]_{ij} - [A_1]_{ji}$, for $1 \leq i, j \leq n$. The goal of this paper is to analyze the geometry of \mathbb{S}_r and, in particular, to answer the following questions:

Q1: Which are the irreducible components of \mathbb{S}_r ?

Q2: Which is the dimension of \mathbb{S}_r ?

One of the motivations to address questions **Q1** and **Q2** is to answer the following question: Given $r < n$, which is the most likely Kronecker canonical form for structured $n \times n$ pencils with rank at most r ? This question is interesting in the context of low-rank perturbations, when one is interested in describing the most-likely (generic) change of the Kronecker canonical form due to low-rank perturbations. In particular, for the unstructured case, as mentioned above, the set of $n \times n$ pencils with rank at most $r < n$ is decomposed into $r + 1$ sets in [7] (which coincide with the irreducible components of the algebraic set of $n \times n$ pencils with rank at most $r < n$ [8]). This allowed to analyze the generic change of the Kronecker canonical form of matrix pencils after low-rank perturbations by looking at the behavior when perturbing with pencils in each of these $r + 1$ sets.

We provide answers to questions **Q1** and **Q2** and, as a consequence, we determine the most-likely (generic) canonical form under congruence of structured matrix pencils with bounded rank, for any of the structures in Table 2.1. Two matrix pencils $A_0 + \lambda A_1$ and $B_0 + \lambda B_1$, are said to be *congruent* if there is some invertible matrix P such that $B_0 + \lambda B_1 = P(A_0 + \lambda A_1)P^\top$. This relation preserves any of the previous structures, that is, if one of $A_0 + \lambda A_1$ or $B_0 + \lambda B_1$ satisfies any of these structures, then the other one does as well. This is no longer true if we replace congruence by strict equivalence, which is the natural relation for unstructured matrix pencils. This is the reason for considering the more restrictive relation of congruence, instead of strict equivalence, in this work. The most remarkable difference with the unstructured case regarding **Q1** is that, while the set of $n \times n$ pencils (non-structured) with rank at most $r < n$ has, as mentioned, $r + 1$ different irreducible components, the set \mathbb{S}_r is irreducible, so it has only one irreducible component, for any of the structures \mathbb{S} in Table 2.1. And regarding the generic canonical form, while in the unstructured case there is no a generic canonical form for $n \times n$ pencils with rank at most r , in the structured case, there is such a generic canonical form.

The paper is organized as follows. In Section 2 we introduce the basic notation and definitions used in the paper. In Section 3 we present a couple of results which are key in the proof of the main results of the paper. These results are provided in Section 4. In particular, Theorem 4.1 shows that the set of $n \times n$ \top -palindromic pencils with rank at most $r < n$ is an irreducible algebraic set, and provides a description of this set as the closure of the congruence orbit of certain pencil given in canonical form. This is, precisely, the generic canonical form for \top -palindromic pencils with rank at most r , as stated in Corollary 4.3. Similar results are presented in Theorem 4.2 for the \top -anti-palindromic structure, and in Theorems 4.4 and 4.5 for the \top -even and \top -odd structures, respectively. In Section 4.3 we compare our results on the dimension of the sets of structured pencils with bounded rank with the dimension of the set of arbitrary $n \times n$ structured pencils (that is, when $r = n$, which allows for full-rank pencils), and also with the set of unstructured pencils. We conclude in Section 5 with a summary of the main contributions of the paper and some indications on further research on the topic.

2. Basic definitions and notation. Throughout the paper we use the following notation. The symbol I_k (or just I , when the size is clear by the context) denotes the $k \times k$ identity matrix. By e_j we denote the j th canonical vector, that is the j th column of the identity matrix, where the size of this matrix depends on the context. The notation A^\top stands for the transpose of the matrix A . The set of vector polynomials with n coordinates (that is, vectors whose n coordinates are polynomials

Structure set \mathbb{S}	Definition $A_0 + \lambda A_1 \in \mathbb{S}$	Notation $\{A_0 + \lambda A_1 \in \mathbb{S} : \text{rank}(A_0 + \lambda A_1) \leq r\}$
\top -palindromic	$A_1^\top = A_0$	Pal_r
\top -anti-palindromic	$A_1^\top = -A_0$	Apal_r
\top -even	$A_0^\top = A_0, A_1^\top = -A_1$	Even_r
\top -odd	$A_0^\top = -A_0, A_1^\top = A_1$	Odd_r

TABLE 2.1

Structured pencils and notation for the sets of structured pencils with $\text{rank} \leq r$

in the variable λ) is denoted by $\mathbb{C}[\lambda]^n$. Given a vector $v(\lambda) \in \mathbb{C}[\lambda]^n$, we denote by $\deg v$ the maximum degree of its coordinates. A *matrix pencil* is a matrix polynomial with degree 1. In other words, a matrix pencil is of the form $A_0 + \lambda A_1$, with $A_0, A_1 \in \mathbb{C}^{m \times n}$. Since we deal only with structured pencils, which must be square, we only consider the case $m = n$. The *reversal* of $A_0 + \lambda A_1$ is the pencil $\text{rev}(A_0 + \lambda A_1) := A_1 + \lambda A_0$. The *rank* of a matrix pencil $A_0 + \lambda A_1$ is the order of the largest non-identically zero minor (considered as a scalar polynomial in λ). In other words, it is the rank of $A_0 + \lambda A_1$ viewed as a matrix in the field of rational functions in the variable λ . This is sometimes referred to in the literature as the *normal rank* (see, for instance, [17]).

We deal with the set of $n \times n$ structured matrix pencils $A_0 + \lambda A_1$ having any of the structures indicated in Table 2.1, and with rank at most $r < n$. This table includes also the abbreviations used in the paper for each of the sets of low-rank structured matrix pencils. The \top -even and \top -odd structures are both gathered under the common denomination \top -*alternating*.

The canonical form under strict equivalence of matrix pencils is the Kronecker canonical form (KCF), which consists of a direct sum of certain (canonical) blocks [19, Ch. XII, §4]. When the pencil enjoys some particular structure, like the ones in Table 2.1, this structure is translated into the KCF in terms of some restrictions in the number and sizes of certain types of blocks. However, the strict equivalence transformations that lead to the KCF do not necessarily preserve the structure. Moreover, the KCF does not fulfill any of the structures in Table 2.1 for general structured pencils. Nonetheless, an equivalent canonical form to the KCF can be achieved using structure-preserving transformations. In particular, for the structures in Table 2.1, these are *congruence transformations*. In other words, given any pencil $S(\lambda)$ fulfilling any of the structures in Table 2.1, there is a nonsingular matrix V such that $VS(\lambda)V^\top$ is in some appropriate canonical form, which depends on the particular structure, and that displays the information contained in the KCF. We will recall later in this section the canonical form for the \top -palindromic structure, which is the only one we need in this paper. For the remaining structures, we refer the reader to [2] and the references therein (see also [23, Cor. 4.3] for the \top -alternating structure). All these canonical forms are, like the KCF, a direct sum of canonical blocks, including some of the following ones.

A *right singular block of order α* is the $\alpha \times (\alpha + 1)$ matrix pencil:

$$L_\alpha(\lambda) := \begin{bmatrix} \lambda & 1 & & & \\ & \ddots & \ddots & & \\ & & \lambda & 1 & \\ & & & & \end{bmatrix}_{\alpha \times (\alpha+1)}.$$

From this block we construct the following four kinds of blocks of size $(2\alpha+1) \times (2\alpha+1)$,

which appear in the canonical form for structured matrix pencils:

$$\begin{aligned}
M_\alpha^\sharp(\lambda) &:= \begin{bmatrix} 0 & L_\alpha(\lambda) \\ \text{rev } L_\alpha(\lambda)^\top & 0 \end{bmatrix}_{(2\alpha+1) \times (2\alpha+1)}, \\
M_\alpha^{-\sharp}(\lambda) &:= \begin{bmatrix} 0 & L_\alpha(\lambda) \\ -\text{rev } L_\alpha(\lambda)^\top & 0 \end{bmatrix}_{(2\alpha+1) \times (2\alpha+1)}, \\
M_\alpha^b(\lambda) &:= \begin{bmatrix} 0 & L_\alpha(\lambda) \\ L_\alpha(-\lambda)^\top & 0 \end{bmatrix}_{(2\alpha+1) \times (2\alpha+1)}, \\
M_\alpha^{-b}(\lambda) &:= \begin{bmatrix} 0 & L_\alpha(\lambda) \\ -L_\alpha(-\lambda)^\top & 0 \end{bmatrix}_{(2\alpha+1) \times (2\alpha+1)}.
\end{aligned}$$

More precisely, the \top -palindromic canonical form contains blocks of type $M_\alpha^\sharp(\lambda)$, the \top -anti-palindromic canonical form contains blocks of type $M_\alpha^{-\sharp}(\lambda)$, the \top -even canonical form contains blocks of type $M_\alpha^b(\lambda)$, and the \top -odd canonical form contains blocks of type $M_\alpha^{-b}(\lambda)$. These are the blocks associated to the *singular spectral structure*. The canonical forms contain also a direct sum of blocks that comprise the *regular spectral structure*, and which are built up from Jordan blocks associated with finite and infinite eigenvalues. A *Jordan block of order k associated with the eigenvalue λ_0* is the following block with size $k \times k$:

$$J_k(\lambda_0) = \begin{bmatrix} \lambda_0 & 1 & & \\ & \ddots & \ddots & \\ & & \lambda_0 & 1 \\ & & & \lambda_0 \end{bmatrix}_{k \times k}.$$

Now we are in the position to state the canonical form for congruence of \top -palindromic matrix pencils. Such a canonical form can be found in [28], but we are using here a different notation. For more details on this canonical form we refer the reader to [3].

THEOREM 2.1. (Canonical form of \top -palindromic pencils). *Any \top -palindromic matrix pencil $L(\lambda)$ is congruent to a direct sum of blocks of the following types:*

- (i) *Blocks of type $M_\alpha^\sharp(\lambda)$.*
- (ii) *Palindromic pairs of Jordan-like blocks with even size associated with $\lambda_0 = -1$:*

$$\begin{bmatrix} 0 & \lambda I_\beta + J_\beta(1) \\ I_\beta + \lambda J_\beta(1)^\top & 0 \end{bmatrix},$$

with β an even number.

- (iii) *Palindromic pairs of Jordan-like blocks with odd size associated with $\lambda_0 = 1$:*

$$\begin{bmatrix} 0 & \lambda I_\gamma + J_\gamma(-1) \\ I_\gamma + \lambda J_\gamma(-1)^\top & 0 \end{bmatrix},$$

with γ an odd number.

- (iv) *Palindromic pairs of Jordan-like blocks associated with $-\lambda_0$ and $-1/\lambda_0$ ($\lambda_0 \neq \pm 1$):*

$$(2.1) \quad \begin{bmatrix} 0 & \lambda I_\delta + J_\delta(\lambda_0) \\ I_\delta + \lambda J_\delta(\lambda_0)^\top & 0 \end{bmatrix}.$$

(v) *Palindromic Jordan-like blocks with even size associated with $\lambda_0 = 1$:*

$$(2.2) \quad \left[\begin{array}{c|c} & \begin{array}{ccc} & & \lambda - 1 \\ & & \lambda - 1 & 1 \\ & & \ddots & \ddots \\ \lambda - 1 & 1 & & \end{array} \\ \hline \begin{array}{ccc} & & 1 - \lambda \\ & 1 - \lambda & \lambda \\ & \ddots & \ddots \\ 1 - \lambda & \lambda & \end{array} & \begin{array}{c} \lambda + 1 \\ \lambda + 1 \end{array} \end{array} \right]_{(2\varepsilon) \times (2\varepsilon)} .$$

(vi) *Palindromic Jordan-like blocks with odd size associated with $\lambda_0 = -1$:*

$$(2.3) \quad \left[\begin{array}{c|c} & \begin{array}{ccc} & & \lambda + 1 \\ & & \ddots & 1 \\ & \lambda + 1 & \ddots & \\ \lambda + 1 & 1 & & \end{array} \\ \hline \begin{array}{ccc} & & \lambda + 1 \\ & 1 + \lambda & \lambda \\ & \ddots & \ddots \\ 1 + \lambda & \lambda & \end{array} & \begin{array}{c} \lambda + 1 \\ \lambda + 1 \end{array} \end{array} \right]_{(2\eta+1) \times (2\eta+1)} .$$

The number of blocks of each type and their particular sizes uniquely depend on $L(\lambda)$ and determine its \top -palindromic canonical form.

The \top -palindromic canonical form in Theorem 2.1 is closely related to the *canonical form for congruence* (CFC) of matrices, so that there is a one-to-one correspondence between blocks in these canonical forms [3, Th. 4]. The only relevant correspondence in our developments is the one between so-called *type 0* blocks, which are Jordan blocks associated with $\lambda_0 = 0$ (see [3, Th. 3]), and blocks of type (i) in Theorem 2.1. More precisely, for each block $J_{2\alpha+1}(0)$ in the CFC of a matrix A there is a block $M_\alpha^\sharp(\lambda)$ in the \top -palindromic canonical form of $A + \lambda A^\top$, and viceversa [3, Th. 4-(ii)]. This correspondence will be used in the proof of Theorem 4.1.

Another relevant notion in this paper is the *orbit under congruence* of a matrix pencil $L(\lambda)$, which is defined as:

$$\mathcal{O}_c(L) = \{V^\top L(\lambda)V : V \text{ nonsingular}\}.$$

The *closure* of this orbit, denoted by $\overline{\mathcal{O}_c(L)}$, is the closure in the standard topology, which is the same as the closure in the Zariski topology [26, Th. 2.33].

DEFINITION 2.2. *Let \mathbb{S}_r be the set of $n \times n$ structured matrix pencils with rank at most r , with \mathbb{S} being any of the structures in Table 2.1. We say that a particular matrix $K_{\mathbb{S}}(\lambda)$ given in structured canonical form is the generic canonical form in \mathbb{S}_r if there is a dense open set of $n \times n$ matrix pencils in \mathbb{S}_r which are congruent to $K_{\mathbb{S}}(\lambda)$.*

In other words, $K_{\mathbb{S}}(\lambda)$ is the generic canonical form in \mathbb{S}_r if $\overline{\mathcal{O}_c(K_{\mathbb{S}})} = \mathbb{S}_r$, since $\mathcal{O}_c(K_{\mathbb{S}})$ is an open set in its closure [21, p. 60].

3. Preliminary results. We first present a decomposition of a given $n \times n$ \top -palindromic matrix pencil with rank at most r as the sum of r rank-1 pencils. This decomposition provides us a constructive way to describe the set of \top -palindromic pencils with bounded rank.

THEOREM 3.1. (Rank-1 decomposition for \top -palindromic pencils). *If $E(\lambda)$ is a \top -palindromic $n \times n$ matrix pencil with $\text{rank } E = r \leq n$, then it can be written as*

$$(3.1) \quad E(\lambda) = \begin{cases} v_1 w_1^\top + \cdots + v_{r/2} w_{r/2}^\top & \text{if } r \text{ is even,} \\ +(\text{rev } w_1) v_1^\top + \cdots + (\text{rev } w_{r/2}) v_{r/2}^\top, & \\ (1 + \lambda) u u^\top + v_1 w_1^\top + \cdots + v_{(r-1)/2} w_{(r-1)/2}^\top & \text{if } r \text{ is odd,} \\ +(\text{rev } w_1) v_1^\top + \cdots + (\text{rev } w_{(r-1)/2}) v_{(r-1)/2}^\top, & \end{cases}$$

where $u, v_1, \dots, v_{\lfloor r/2 \rfloor} \in \mathbb{C}^n$ and $w_1, \dots, w_{\lfloor r/2 \rfloor} \in \mathbb{C}[\lambda]^n$ with $\deg w_i \leq 1$, for $i = 1, \dots, \lfloor r/2 \rfloor$.

Proof. Let us assume that the result is true for any \top -palindromic pencil being in \top -palindromic canonical form as in Theorem 2.1. Let $E(\lambda)$ be an arbitrary \top -palindromic pencil. By Theorem 2.1, there is some invertible matrix P such that $PE(\lambda)P^\top = K_E(\lambda)$, with $K_E(\lambda)$ being in \top -palindromic canonical form. Then $K_E(\lambda)$ is of the form (3.1). Now, by setting

$$\tilde{v}_i = P^{-1}v_i, \quad \tilde{w}_i = P^{-1}w_i, \quad \tilde{u} = P^{-1}u,$$

with v_i, w_i, u as in (3.1), we arrive at a decomposition like (3.1) for $E(\lambda)$, with \tilde{v}_i, \tilde{w}_i , and \tilde{u} instead of v_i, w_i, u , respectively (note that, since P^{-1} is invertible, $\text{rev}(P^{-1}v) = P^{-1} \text{rev } v$ and $\deg(P^{-1}v) = \deg v$, for any $v \in \mathbb{C}[\lambda]^n$).

Therefore, we may assume that $E(\lambda)$ is given in \top -palindromic canonical form. Then, it is a direct sum of blocks of types (i)–(vi) in Theorem 2.1. We are going to show that each of these blocks can be decomposed as a sum of rank-1 pencils in such a way that the whole direct sum is of the form (3.1). Let us show such a decomposition for each type of canonical blocks.

- A block of type (i) can be written as

$$M_\alpha^\#(\lambda) = e_1(\lambda e_{\alpha+1} + e_{\alpha+2})^\top + \cdots + e_\alpha(\lambda e_{2\alpha} + e_{2\alpha+1})^\top \\ + (e_{\alpha+1} + \lambda e_{\alpha+1}) e_1^\top + \cdots + (e_{2\alpha} + \lambda e_{2\alpha+1}) e_\alpha^\top.$$

- A block of type (iv) like (2.1) can be written as

$$e_1((\lambda + \lambda_0)e_{\delta+1} + e_{\delta+2})^\top + \cdots + e_{\delta-1}((\lambda + \lambda_0)e_{2\delta-1} + e_{2\delta})^\top + e_\delta((\lambda + \lambda_0)e_{2\delta})^\top \\ + (1 + \lambda\lambda_0)e_{\delta+1} + \lambda e_{\delta+2}) e_1^\top + \cdots + ((1 + \lambda\lambda_0)e_{2\delta+1} + \lambda e_{2\delta}) e_{\delta-1}^\top + ((1 + \lambda\lambda_0)e_{2\delta}) e_\delta^\top.$$

- A block of type (v) like (2.2) can be decomposed as

$$e_{2\varepsilon}((1 - \lambda)e_1 + \lambda e_2)^\top + \cdots + e_{\varepsilon+2}((1 - \lambda)e_{\varepsilon-1} + \lambda e_\varepsilon)^\top \\ + e_{\varepsilon+1}((1 - \lambda)e_\varepsilon + \lambda e_{\varepsilon+1})^\top \\ + ((\lambda - 1)e_1 + e_2) e_{2\varepsilon}^\top + \cdots + ((\lambda - 1)e_{\varepsilon-1} + e_\varepsilon) e_{\varepsilon+2}^\top \\ + ((\lambda - 1)e_\varepsilon + e_{\varepsilon+1}) e_{\varepsilon+1}^\top.$$

- A block of type (vi) like (2.3) can be decomposed as

$$e_{2\eta+1}((1 + \lambda)e_1 + \lambda e_2)^\top + \cdots + e_{\eta+2}((1 + \lambda)e_\eta + \lambda e_{\eta+1})^\top \\ + ((\lambda + 1)e_1 + e_2) e_{2\eta+1}^\top + \cdots + ((\lambda + 1)e_\eta + e_{\eta+1}) e_{\eta+2}^\top \\ + (\lambda + 1)e_{\eta+1} e_{\eta+1}^\top.$$

- For blocks of types (ii) and (iii) the decomposition is similar to the one for blocks of type (iv), replacing $\lambda_0 = \pm 1$, and δ by either β or γ .

Now, joining up in a direct sum all blocks in the \top -palindromic canonical form of $E(\lambda)$, and padding up with zeroes in all canonical vectors, for each block in the previous rank-1 decompositions, in all positions corresponding to the remaining blocks, we end up with a rank-1 decomposition of the form

$$v_1 w_1^\top + \cdots + v_s w_s^\top + (\text{rev } w_1) v_1^\top + \cdots + (\text{rev } w_s) v_s^\top + (1 + \lambda)(u_1 u_1^\top + \cdots + u_t u_t^\top),$$

with $2s + t = r$, since the rank of $E(\lambda)$ is the sum of the ranks of all canonical blocks. The summands of the form $(1 + \lambda)u_i u_i^\top$ in the previous decomposition come from blocks of type (vi). Given a pair of vectors $u, \tilde{u} \in \mathbb{C}^n$, we can write

$$(1 + \lambda)(uu^\top + \tilde{u}\tilde{u}^\top) = vw^\top + (\text{rev } w)v^\top,$$

with $v = u + i\tilde{u}$, $w = \frac{1+\lambda}{2}(u - i\tilde{u})$ (where i denotes the imaginary unit). Therefore, if r is even, we can gather these summands in couples to get a decomposition like in the first expression of (3.1). However, when r is odd, one of the summands remains unpaired, and we arrive at the second expression in (3.1). This proves the result. \square

The following result, which is closely connected to Theorem 3.1, deals with the set of $n \times n$ matrices that can be decomposed as a sum of rank-1 matrices in a specific way. It will be key in computing the dimension of the set of \top -palindromic pencils with bounded rank, which will be key in turn in the proof of Theorem 4.1.

PROPOSITION 3.2. *Let s, n be two integers with $0 < s \leq n$. Let us define the following sets of $n \times n$ matrices with complex entries:*

$$(3.2) \quad \mathcal{M}_s = \left\{ u_1 v_1^\top + \cdots + u_s v_s^\top + w_1 u_1^\top + \cdots + w_s u_s^\top : \begin{array}{l} u_i, v_i, w_i \in \mathbb{C}^n, \\ \text{for } i = 1, \dots, s \end{array} \right\}, \text{ and}$$

$$(3.3) \quad \widetilde{\mathcal{M}}_s = \left\{ uu^\top + u_1 v_1^\top + \cdots + u_s v_s^\top + w_1 u_1^\top + \cdots + w_s u_s^\top : \begin{array}{l} u, u_i, v_i, w_i \in \mathbb{C}^n, \\ \text{for } i = 1, \dots, s \end{array} \right\}.$$

Then \mathcal{M}_s and $\widetilde{\mathcal{M}}_s$ are algebraic sets and

- (a) $\dim \mathcal{M}_s \leq s(3n - 2s)$,
- (b) $\dim \widetilde{\mathcal{M}}_s \leq s(3n - 2s - 1) + n$.

Proof. Let us define the maps

$$\begin{aligned} \Phi_1 : \quad & \mathbb{C}^{sn} \times \mathbb{C}^{sn} \times \mathbb{C}^{sn} & \longrightarrow & \mathbb{C}^{n \times n} \\ & (u_1, \dots, u_s; v_1, \dots, v_s; w_1, \dots, w_s) & \longmapsto & \begin{array}{l} u_1 v_1^\top + \cdots + u_s v_s^\top \\ + w_1 u_1^\top + \cdots + w_s u_s^\top, \end{array} \end{aligned}$$

and

$$\begin{aligned} \Phi_2 : \quad & \mathbb{C}^n \times \mathbb{C}^{sn} \times \mathbb{C}^{sn} \times \mathbb{C}^{sn} & \longrightarrow & \mathbb{C}^{n \times n} \\ & (u; u_1, \dots, u_s; v_1, \dots, v_s; w_1, \dots, w_s) & \longmapsto & \begin{array}{l} uu^\top + u_1 v_1^\top + \cdots + u_s v_s^\top \\ + w_1 u_1^\top + \cdots + w_s u_s^\top. \end{array} \end{aligned}$$

Both Φ_1 and Φ_2 are regular maps, since they are defined by homogeneous polynomials of the same degree (they are all quadratic polynomials in the coordinates of u and u_i, v_i, w_i , for $i = 1, \dots, s$). The sets \mathcal{M}_s and $\widetilde{\mathcal{M}}_s$ are the images of, respectively, Φ_1 and Φ_2 . Since the image of an algebraic set under a regular map is closed [29, §5.2, Th. 2], we conclude that both \mathcal{M}_s and $\widetilde{\mathcal{M}}_s$ are algebraic sets.

In order to prove the second part of the statement, we look at the tangent space of each \mathcal{M}_s and $\widetilde{\mathcal{M}}_s$ at a general point. Let us first consider the set \mathcal{M}_s . The tangent

space is spanned by the $3sn$ vectors obtained by taking partial derivatives in Φ_1 , namely the following $3sn$ matrices with size $n \times n$:

$$(3.4) \quad \begin{aligned} e_j v_i^\top + w_i e_j^\top, & \quad i = 1, \dots, s, \quad j = 1, \dots, n, \\ u_i e_j^\top, & \quad i = 1, \dots, s, \quad j = 1, \dots, n, \\ e_j u_i^\top, & \quad i = 1, \dots, s, \quad j = 1, \dots, n. \end{aligned}$$

Let us write the matrices in (3.4) as vectors in \mathbb{C}^{n^2} using the vec operator [20, Def. 4.2.9], so that they become

$$(3.5) \quad \begin{aligned} v_i \otimes e_j + e_j \otimes w_i, & \quad i = 1, \dots, s, \quad j = 1, \dots, n, \\ e_j \otimes u_i, & \quad i = 1, \dots, s, \quad j = 1, \dots, n, \\ u_i \otimes e_j, & \quad i = 1, \dots, s, \quad j = 1, \dots, n, \end{aligned}$$

where \otimes denotes the Kronecker product [20, Def. 4.2.1]. We are going to see that the set of vectors (3.4) contains at least $2s^2$ linearly dependent vectors. A null linear combination of the vectors in (3.5) can be written as

$$(3.6) \quad \sum_{i,j} [x_{ji}(v_i \otimes e_j + e_j \otimes w_i) + y_{ji}(e_j \otimes u_i) + z_{ji}(u_i \otimes e_j)] = M \begin{bmatrix} \text{vec}(X) \\ \text{vec}(Y) \\ \text{vec}(Z) \end{bmatrix} = 0,$$

with $X = [x_{ij}]$, $Y = [y_{ij}]$, $Z = [z_{ij}]$, for $1 \leq i \leq s$, $1 \leq j \leq n$, and $M = [M_1 \mid M_2 \mid M_3]$, where M_1 contains all columns of the form $v_i \otimes e_j + e_j \otimes w_i$, M_2 contains all columns of the form $e_j \otimes u_i$, and M_3 contains all columns of the form $u_i \otimes e_j$ in the left-hand side of (3.6), and they all are ordered in the lexicographic order of the pairs (i, j) .

The left-hand side of (3.6) can be written as

$$\sum_{i=1}^s \left[v_i \otimes \left(\sum_{j=1}^n x_{ji} e_j \right) + \left(\sum_{j=1}^n x_{ji} e_j \right) \otimes w_i + \left(\sum_{j=1}^n y_{ji} e_j \right) \otimes u_i + u_i \otimes \left(\sum_{j=1}^n z_{ji} e_j \right) \right].$$

If we denote the columns of X , Y , and Z , respectively, by

$$\begin{aligned} \bar{x}_i &:= \text{Col}_i X = \sum_{j=1}^n x_{ji} e_j, & i = 1, \dots, s, \\ \bar{y}_i &:= \text{Col}_i Y = \sum_{j=1}^n y_{ji} e_j, & i = 1, \dots, s, \\ \bar{z}_i &:= \text{Col}_i Z = \sum_{j=1}^n z_{ji} e_j, & i = 1, \dots, s, \end{aligned}$$

then (3.6) can be written as

$$(3.7) \quad \sum_{i=1}^s [v_i \otimes \bar{x}_i + \bar{x}_i \otimes w_i + \bar{y}_i \otimes u_i + u_i \otimes \bar{z}_i] = 0.$$

Now, we can construct $2s^2$ different solutions to (3.6) as follows. Given a pair (i_0, j_0) with $1 \leq i_0, j_0 \leq s$, we set:

$$(3.8) \quad \begin{aligned} s_1(i_0, j_0) &= e_{s+i_0} \otimes u_{j_0} - e_{2s+j_0} \otimes u_{i_0}, & \text{and} \\ s_2(i_0, j_0) &= e_{i_0} \otimes u_{j_0} - e_{s+j_0} \otimes v_{i_0} - e_{2s+j_0} \otimes w_{i_0}, \end{aligned}$$

where the canonical vectors in (3.8) belong to \mathbb{C}^{3s} . It is straightforward to check that, for a fixed pair (i_0, j_0) with $1 \leq i_0, j_0 \leq s$, the vector $s_1(i_0, j_0)$ corresponds to replacing $\bar{y}_{i_0} = u_{j_0}$, $\bar{z}_{j_0} = -u_{i_0}$, and the remaining vectors $\bar{x}_i = \bar{y}_i = \bar{z}_i = 0$ in (3.7). Similarly,

the vector $s_2(i_0, j_0)$ corresponds to replacing $\bar{x}_{i_0} = u_{j_0}, \bar{y}_{j_0} = -v_{i_0}, \bar{z}_{j_0} = -w_{i_0}$, and the remaining vectors $\bar{x}_i = \bar{y}_i = \bar{z}_i = 0$ in (3.7). Then, $s_1(i_0, j_0)$ and $s_2(i_0, j_0)$ give $2s^2$ different solutions of (3.6) for general vectors u_i, v_i, w_i . Now, let us prove that, for general vectors u_i, v_i, w_i , these solutions are linearly independent.

Let us assume that there is a null linear combination of the solutions

$$(3.9) \quad \sum_{i,j=1}^s \alpha_{ij} s_1(i, j) + \sum_{i,j=1}^s \beta_{ij} s_2(i, j) = 0,$$

with $\alpha_{ij}, \beta_{ij} \in \mathbb{C}$, where we have replaced (i_0, j_0) by (i, j) for simplicity. Then, replacing (3.8) into this expression, we arrive at

$$(3.10) \quad \sum_{i,j=1}^s \alpha_{ij} (e_{s+i} \otimes u_j - e_{2s+j} \otimes u_i) + \sum_{i,j=1}^s \beta_{ij} (e_i \otimes u_j - e_{s+j} \otimes v_i - e_{2s+j} \otimes w_i) = 0.$$

Looking at the summands in the left-hand side of (3.10) whose first vector in the Kronecker product is of the form e_k , for $1 \leq k \leq s$, and equating to zero, we arrive at

$$(3.11) \quad \begin{aligned} \beta_{11} u_1 + \cdots + \beta_{1s} u_s &= 0, \\ \vdots & \\ \beta_{s1} u_1 + \cdots + \beta_{ss} u_s &= 0. \end{aligned}$$

For a general point in Pal_r , the set $\{u_1, \dots, u_s\}$ is linearly independent, since $s \leq n$, by hypothesis. Therefore, (3.11) implies that $\beta_{ij} = 0$, for all $1 \leq i, j \leq s$. Now, looking at the summands in (3.10) whose first vector in the Kronecker product is of the form e_k , with $s+1 \leq k \leq 2s$, and equating to zero, we arrive at

$$(3.12) \quad \begin{aligned} \alpha_{11} u_1 + \cdots + \alpha_{1s} u_s &= 0, \\ \vdots & \\ \alpha_{s1} u_1 + \cdots + \alpha_{ss} u_s &= 0. \end{aligned}$$

Again, the set $\{u_1, \dots, u_s\}$ is linearly independent for a general point in Pal_r , so (3.12) implies that $\alpha_{ij} = 0$, for all $1 \leq i, j \leq s$. Therefore, the only null linear combination (3.9) for a general point in Pal_r is the one with $\alpha_{ij} = \beta_{ij} = 0$, for all $1 \leq i, j \leq s$, which implies that $s_1(i, j)$ and $s_2(i, j)$, for $1 \leq i, j \leq s$, are linearly independent, as wanted. As a consequence, $\dim \mathcal{M}_s \leq 3ns - 2s^2 = s(3n - 2s)$, as claimed.

Now, let us address the proof of the second part of the statement for the set $\widetilde{\mathcal{M}}_s$. The tangent space at a given point is the linear space spanned by the following $3sn + n$ vectors, obtained from the partial derivatives of Φ_2 :

$$\begin{aligned} e_j v_i^\top + w_i e_j^\top, & \quad i = 1, \dots, s, \quad j = 1, \dots, n, \\ u_i e_j^\top, & \quad i = 1, \dots, s, \quad j = 1, \dots, n, \\ e_j u_i^\top, & \quad i = 1, \dots, s, \quad j = 1, \dots, n, \\ e_j u^\top + u e_j^\top, & \quad j = 1, \dots, n, \end{aligned}$$

and, applying again the vec operator, these vectors become

$$(3.13) \quad \begin{aligned} v_i \otimes e_j + e_j \otimes w_i, & \quad i = 1, \dots, s, \quad j = 1, \dots, n, \\ e_j \otimes u_i, & \quad i = 1, \dots, s, \quad j = 1, \dots, n, \\ u_i \otimes e_j, & \quad i = 1, \dots, s, \quad j = 1, \dots, n, \\ u \otimes e_j + e_j \otimes u, & \quad j = 1, \dots, n. \end{aligned}$$

A null linear combination of the vectors in (3.13) is of the form

$$(3.14) \quad \sum_{i,j} [x_{ji}(v_i \otimes e_j + e_j \otimes w_i) + y_{ji}(e_j \otimes u_i) + z_{ji}(u_i \otimes e_j) + t_j(u \otimes e_j + e_j \otimes u)] \\ = \widetilde{M} \begin{bmatrix} \text{vec}(X) \\ \text{vec}(Y) \\ \text{vec}(Z) \\ \text{vec}(T) \end{bmatrix} = 0,$$

with $X = [x_{ij}]$, $Y = [y_{ij}]$, $Z = [z_{ij}]$, $T = [t_i]$, for $1 \leq i \leq s$, $1 \leq j \leq n$, and $M = [M_1 \mid M_2 \mid M_3 \mid M_4]$, where M_1 contains all columns of the form $v_i \otimes e_j + e_j \otimes w_i$, M_2 contains all columns of the form $e_j \otimes u_i$, M_3 contains all columns of the form $u_i \otimes e_j$, and M_4 contains all columns of the form $u \otimes e_i + e_i \otimes u$ in the left-hand side of (3.14), and they all are ordered in the lexicographic order of the pairs (i, j) . With the same notation $\bar{x}_i, \bar{y}_i, \bar{z}_i$ for the columns of X, Y, Z as before, together with $\bar{t} = \sum_{j=1}^n t_j e_j$, Equation (3.14) is equivalent to

$$(3.15) \quad \sum_{i=1}^s [v_i \otimes \bar{x}_i + \bar{x}_i \otimes w_i + \bar{y}_i \otimes u_i + u_i \otimes \bar{z}_i] + u \otimes \bar{t} + \bar{t} \otimes u = 0.$$

Now, for each (i_0, j_0) , with $1 \leq i_0, j_0 \leq s$, we define $s_1(i_0, j_0)$ and $s_2(i_0, j_0)$ as in (3.8), with the only difference that now the canonical vectors belong to \mathbb{C}^{3s+1} . We also set, for each $1 \leq i_0 \leq s$,

$$(3.16) \quad s_3(i_0) = -e_{s+i_0} \otimes u_{i_0} - e_{2s+i_0} \otimes u_{i_0} + e_{3s+1} \otimes u_{i_0},$$

where, again, the canonical vectors in (3.16) belong to \mathbb{C}^{3s+1} . The vectors (3.8) are solutions of (3.15), for the same reason as in the preceding case. The n vectors $s_3(i)$ in (3.16) correspond to replacing $\bar{y}_{i_0} = \bar{z}_{i_0} = -u_{i_0}$, $\bar{t} = u_{i_0}$, and $\bar{x}_i, \bar{y}_i, \bar{z}_i = 0$, for $i \neq i_0$ in (3.14). Then, $s_3(i)$ is also a solution of (3.15), for $1 \leq i \leq s$. It remains to prove that $s_1(i, j)$, $s_2(i, j)$, and $s_3(i)$ are linearly independent for a general vector in Pal_r . This follows similar arguments to the ones for \mathcal{M}_s . In particular, if

$$(3.17) \quad \sum_{i,j=1}^s \alpha_{ij} s_1(i, j) + \sum_{i,j=1}^s \beta_{ij} s_2(i, j) + \sum_{i=1}^s \gamma_i s_3(i) = 0$$

is a null linear combination of $s_1(i, j)$, $s_2(i, j)$, and $s_3(i)$, then replacing (3.8) and (3.16) in (3.17) we arrive at

$$(3.18) \quad \sum_{i,j=1}^s \alpha_{ij} (e_{s+i} \otimes u_i - e_{2s+j} \otimes u_j) + \sum_{i,j=1}^s \beta_{ij} (e_i \otimes u_j - e_{s+j} \otimes v_i - e_{2s+j} \otimes w_i) \\ + \sum_{i=1}^s \gamma_i (-e_{s+i} \otimes u_i - e_{2s+i} \otimes u_i + e_{3s+1} \otimes u_i) = 0.$$

Looking at the summands whose first term in the Kronecker product is of the form e_{3s+1} we arrive at $\gamma_1 u_1 + \dots + \gamma_s u_s = 0$ and this implies, provided that $\{u_1, \dots, u_s\}$ is linearly independent, that $\gamma_1 = \dots = \gamma_s = 0$. Then, looking again at the terms whose first vector in the Kronecker product is of the form e_k , with $1 \leq k \leq s$, we conclude that, as long as $\{u_1, \dots, u_s\}$ is linearly independent, $\beta_{ij} = 0$, for all $1 \leq i, j \leq s$. Finally, looking at the terms whose first vector in the Kronecker product is of the form e_k , with $s+1 \leq k \leq 2s$, we get $\alpha_{ij} = 0$, for all $1 \leq i, j \leq s$. This implies that the $2s^2 + s$ solutions $s_1(i, j)$, $s_2(i, j)$, and $s_3(j)$ are linearly independent, so $\dim \widetilde{M}_s \leq 3sn + n - 2s^2 - s = s(3n - 2s - 1) + n$, as wanted. \square

4. Main results. The main results in Sections 4.1 and 4.2 are the analogues of those in the recent paper [12], for skew-symmetric matrix pencils. The proof for the skew-symmetric structure in that paper is based on the fact that a given skew-symmetric pencil $S_1(\lambda)$ is in the closure of the congruence orbit of another skew-symmetric pencil $S_2(\lambda)$ if and only if $S_1(\lambda)$ is in the closure of the strict equivalence orbit of $S_2(\lambda)$. In other words, if there is a sequence of pencils strictly equivalent to $S_2(\lambda)$ which converges to $S_1(\lambda)$, then there is also a sequence of pencils which are congruent to $S_2(\lambda)$ and that converges to $S_1(\lambda)$. This is a very strong result from [13], and it is not yet known whether an analogous result is true or not for \mathbb{T} -palindromic or \mathbb{T} -alternating structures. Therefore, a relevant part of the proof of Theorem 4.1, which is the main result in Section 4.1, follows a completely different technique to the ones in [12], relying on Proposition 3.2.

We analyze separately the following structures: (i) \mathbb{T} -palindromic and \mathbb{T} -anti-palindromic structures (Section 4.1), and (ii) \mathbb{T} -alternating structures (Section 4.2). The \mathbb{T} -palindromic and \mathbb{T} -anti-palindromic structures are related to each other by the elementary change of variables $\lambda \mapsto -\lambda$, so the results for one of these structures are directly extended to the other one. Similarly, the \mathbb{T} -even and \mathbb{T} -odd structures are related by reversing the order of the coefficients A_0 and A_1 , so it is again enough to analyze just one of them. The \mathbb{T} -palindromic and the \mathbb{T} -alternating structures are also related by particular cases of Möbius transformations (known as *Cayley transformations*). Using these transformations, the results for \mathbb{T} -palindromic pencils can be easily translated to \mathbb{T} -alternating pencils as well.

4.1. \mathbb{T} -palindromic and \mathbb{T} -anti-palindromic pencils. Our main results in this section show that the sets of \mathbb{T} -palindromic and \mathbb{T} -anti-palindromic matrix pencils with bounded (deficient) rank are irreducible, and provide the dimension of these sets. They also provide the generic canonical form of these pencils. We start with the \mathbb{T} -palindromic structure.

THEOREM 4.1. (The set of \mathbb{T} -palindromic pencils with bounded rank). *Let r be an integer with $0 \leq r < n$. The set Pal_r is an irreducible algebraic set with dimension*

$$\dim \text{Pal}_r = \begin{cases} \frac{r}{2} \cdot (3n - r), & \text{if } r \text{ is even,} \\ \frac{r-1}{2} \cdot (3n - r) + n, & \text{if } r \text{ is odd.} \end{cases}$$

Moreover, if r is even, then Pal_r is the closure of the congruence orbit of the pencil

$$(4.1) \quad K_P^e(\lambda) := \text{diag}(\overbrace{M_{\alpha+1}^\sharp(\lambda), \dots, M_{\alpha+1}^\sharp(\lambda)}^s, \overbrace{M_\alpha^\sharp(\lambda), \dots, M_\alpha^\sharp(\lambda)}^{n-r-s}),$$

where $r/2 = (n-r)\alpha + s$ is the Euclidean division of $r/2$ by $n-r$. If r is odd, then Pal_r is the closure of the congruence orbit of

$$(4.2) \quad K_P^o(\lambda) := \text{diag}(1 + \lambda, K_P^e(\lambda)),$$

with $K_P^e(\lambda)$ as in (4.1), but now $(r-1)/2 = (n-r)\alpha + s$ is the Euclidean division of $(r-1)/2$ by $n-r$.

Proof. Let us first consider the case r even. The codimension of the orbit of $K_P^e(\lambda)$ in (4.1) can be computed using the formula in [6, Th. 2] and the relationship between $\text{KCF}(A + \lambda A^\top)$ and the CFC of A provided in [3, Th. 4] (see the paragraph right after Theorem 2.1). In particular, if $K_P^e(\lambda) = A + \lambda A^\top$, then

$$\text{CFC}(A) = \text{diag}(\underbrace{J_{2\alpha+3}(0), \dots, J_{2\alpha+3}(0)}_s, \underbrace{J_{2\alpha+1}(0), \dots, J_{2\alpha+1}(0)}_{n-r-s}).$$

Now, applying [6, Th. 2], the codimension of $K_P^e(\lambda)$ is $c_{K_P^e} = c_0 + c_{00}$. The quantity c_0 is the ‘‘codimension’’ of individual blocks $J_k(0)$, and is obtained by adding up $\lceil k/2 \rceil$, for each block $J_k(0)$, whereas c_{00} is due to the ‘‘interactions’’ between two different blocks, taking each pair $(J_k(0), J_\ell(0))$ with $k \leq \ell$ only once, and is equal to: (a) k , if k is even, (b) ℓ , if k is odd and $k \neq \ell$, and (c) $k + 1$ if k is odd and $k = \ell$. In particular, for the blocks in $\text{CFC}(A)$ above,

$$(4.3) \quad c_0 = \sum_{i=1}^s \left\lceil \frac{2\alpha + 3}{2} \right\rceil + \sum_{i=1}^{n-r-s} \left\lceil \frac{2\alpha + 1}{2} \right\rceil = s(\alpha + 2) + (n - r - s)(\alpha + 1) = n - \frac{r}{2},$$

and

$$(4.4) \quad c_{00} = \binom{s}{2} (2\alpha + 4) + \binom{n-r-s}{2} (2\alpha + 2) + s(n - r - s)(2\alpha + 1).$$

After some manipulations in (4.4) we arrive at

$$c_{00} = (n - r - 1) \left(n - \frac{r}{2} \right).$$

Now, adding up, we get $c_{K_P^e} = c_0 + c_{00} = (n - r/2)(n - r)$. Then, the dimension of the congruence orbit of $K_P^e(\lambda)$ is

$$\dim \mathcal{O}_c(K_P^e) = n^2 - c_{K_P^e} = \frac{r}{2}(3n - r).$$

Now, since $\mathcal{O}_c(K_P^e) \subseteq \text{Pal}_r$ and Pal_r is an algebraic (hence closed) set, it follows that $\overline{\mathcal{O}_c(K_P^e)} \subseteq \text{Pal}_r$. In order to prove that the inclusion is an identity, it suffices to see that the dimension of Pal_r is, at most, $\frac{r}{2}(3n - r)$, and that Pal_r is irreducible.

By decomposing $w_i = w_{i0} + \lambda w_{i1}$ in (3.1), for $i = 1, \dots, r/2$, any \top -palindromic pencil $L(\lambda)$ with rank at most r can be written as

$$L(\lambda) = v_1 w_{10}^\top + \dots + v_{r/2} w_{r/2,0}^\top + w_{11} v_1^\top + \dots + w_{r/2,1} v_{r/2}^\top + \lambda (w_{10} v_1^\top + \dots + w_{r/2,0} v_{r/2}^\top + v_1 w_{11}^\top + \dots + v_{r/2} w_{r/2,1}^\top).$$

Then, the pencil $L(\lambda)$ is uniquely determined by its trailing coefficient. In other words, the set Pal_r is in one-to-one correspondence with the set of matrices that can be written in the form $u_1 v_1^\top + \dots + u_{r/2} v_{r/2}^\top + w_1 u_1^\top + \dots + w_{r/2} u_{r/2}^\top$. Proposition 3.2 (a) with $s = r/2$ guarantees that the dimension of this set is at most $\frac{r}{2}(3n - r)$.

To prove that Pal_r is irreducible we proceed as follows. Pal_r is the image of the following (polynomial) map:

$$\begin{aligned} \Phi : \quad \mathbb{C}^{\frac{3rn}{2}} &\longrightarrow \mathbb{C}^{n \times n} \times \mathbb{C}^{n \times n} \\ (v_1, \dots, v_{r/2}; & \quad (v_1 w_{10}^\top + \dots + v_{r/2} w_{r/2,0}^\top + w_{11} v_1^\top + \dots + w_{r/2,1} v_{r/2}^\top, \\ w_{10}, \dots, w_{r/2,0}; & \quad w_{10} v_1^\top + \dots + w_{r/2,0} v_{r/2}^\top + v_1 w_{11}^\top + \dots + v_{r/2} w_{r/2,1}^\top). \\ w_{11}, \dots, w_{r/2,1}) & \end{aligned}$$

Then, assume $\text{Pal}_r = \Phi(\mathbb{C}^{\frac{3rn}{2}}) = X \cup Y$, with X, Y being algebraic sets, which is equivalent to $\mathbb{C}^{\frac{3rn}{2}} = \Phi^{-1}(X) \cup \Phi^{-1}(Y)$. Since $\mathbb{C}^{\frac{3rn}{2}}$ is irreducible and both $\Phi^{-1}(X)$ and $\Phi^{-1}(Y)$ are algebraic sets, this implies that either $\mathbb{C}^{\frac{3rn}{2}} = \Phi^{-1}(X)$ or $\mathbb{C}^{\frac{3rn}{2}} = \Phi^{-1}(Y)$, which in turn implies either $\Phi(\mathbb{C}^{\frac{3rn}{2}}) = X$ or $\Phi(\mathbb{C}^{\frac{3rn}{2}}) = Y$. As a consequence, $\Phi(\mathbb{C}^{\frac{3rn}{2}}) = \text{Pal}_r$ is irreducible, and the proof for the case r even is complete.

Now, let us consider the case r odd. In this case (see [3, Th. 4] or the paragraph right after Theorem 2.1),

$$\text{CFC}(A) = \text{diag}(\underbrace{J_{2\alpha+3}(0), \dots, J_{2\alpha+3}(0)}_s, \underbrace{J_{2\alpha+1}(0), \dots, J_{2\alpha+1}(0)}_{n-r-s}, 1).$$

Therefore, the codimension of $\mathcal{O}_c(K_P^o)$ is $c_{K_P^o} = c_0 + c_{00} + c_{01}$, with c_0 and c_{00} as in (4.3) and (4.4), respectively. The term c_{01} is due to the presence of the last block equal to 1 in $\text{CFC}(A)$, which is a so-called *type I* block [3, Th. 3]. For a given matrix A , the term c_{01} is equal to the product of the number of type 0 blocks in $\text{CFC}(A)$ (that is, blocks of the form $J_k(0)$) and the sum of the sizes of all type I blocks in $\text{CFC}(A)$ (see [6, Th. 2]). In the case of A above, it is equal to $c_{01} = n - r$. Now, however, $c_0 = n - (r + 1)/2$, and with similar manipulations as for the r even case, we can get $c_{00} = (n - r - 1)(n - (r + 1)/2)$. Adding up

$$c_{K_P^o} = c_0 + c_{00} + c_{01} = (n - r) \left(n - \frac{r - 1}{2} \right).$$

Then, the dimension of the congruence orbit of $K_P^o(\lambda)$ is

$$\dim \mathcal{O}_c(K_P^o) = n^2 - c_{K_P^o} = \frac{r - 1}{2} \cdot (3n - r) + n.$$

Again, it remains to prove that the dimension of Pal_r is, at most, $\frac{r - 1}{2} \cdot (3n - r) + n$ and that Pal_r is irreducible. Writing again $w_i = w_{i0} + \lambda w_{i1}$ in (3.1), for $i = 1, \dots, (r - 1)/2$, any \top -palindromic matrix pencil $L(\lambda)$ with rank at most r can be written as

$$\begin{aligned} L(\lambda) = & uu^\top + v_{10}w_{10}^\top + \dots + v_{r/2}w_{r/2,0}^\top + w_{11}v_1^\top + \dots + w_{r/2,1}v_{r/2}^\top \\ & + \lambda(w_{10}v_1^\top + \dots + w_{r/2,0}v_{r/2}^\top + v_1w_{11}^\top + \dots + v_{r/2}w_{r/2,1}^\top). \end{aligned}$$

As before, the pencil $L(\lambda)$ is uniquely determined by its trailing coefficient. In other words, the set Pal_r is in one-to-one correspondence with the set of matrices that can be written in the form $uu^\top + u_1v_1^\top + \dots + u_{(r-1)/2}v_{(r-1)/2}^\top + w_1u_1^\top + \dots + w_{(r-1)/2}u_{(r-1)/2}^\top$. Proposition 3.2 (b) with $s = (r - 1)/2$ guarantees that the dimension of this set is at most $\frac{r-1}{2}(3n - r) + n$, as wanted.

The proof of the irreducibility of Pal_r in this case follows the same arguments as for the r even case. \square

Now we state the counterpart of Theorem 4.1 for \top -anti-palindromic pencils.

THEOREM 4.2. (The set of \top -anti-palindromic pencils with bounded rank). *Let r be an integer with $0 \leq r < n$. The set Apal_r is an irreducible algebraic set with dimension*

$$\dim \text{Apal}_r = \begin{cases} \frac{r}{2} \cdot (3n - r), & \text{if } r \text{ is even,} \\ \frac{r - 1}{2} \cdot (3n - r) + n, & \text{if } r \text{ is odd.} \end{cases}$$

Moreover, if r is even, then Apal_r is the closure of the congruence orbit of the pencil

$$(4.5) \quad K_A^e(\lambda) := \text{diag}(\underbrace{M_{\alpha+1}^{-\#}(\lambda), \dots, M_{\alpha+1}^{-\#}(\lambda)}_s, \underbrace{M_{\alpha}^{-\#}(\lambda), \dots, M_{\alpha}^{-\#}(\lambda)}_{n-r-s}),$$

where $r/2 = (n-r)\alpha + s$ is the Euclidean division of $r/2$ by $n-r$. If r is odd, then Apal_r is the closure of the congruence orbit of

$$(4.6) \quad K_A^o(\lambda) := \text{diag}(1 - \lambda, K_A^e(\lambda)),$$

with $K_A^e(\lambda)$ as in (4.5), but now $(r-1)/2 = (n-r)\alpha + s$ is the Euclidean division of $(r-1)/2$ by $n-r$.

Proof. The result is an immediate consequence of Theorem 4.1, since a matrix pencil $P(\lambda)$ is \top -anti-palindromic if and only if $P(-\lambda)$ is \top -palindromic. \square

Theorems 4.1 and 4.2 give the generic canonical form of \top -palindromic and \top -anti-palindromic $n \times n$ pencils with rank at most r .

COROLLARY 4.3. (Generic canonical form of \top -palindromic and \top -anti-palindromic pencils with bounded rank). *Let $0 \leq r < n$. The generic canonical structure of $n \times n$ \top -palindromic (respectively, \top -anti-palindromic) $n \times n$ matrix pencils with rank at most r is (4.1) (resp., (4.5)) if r is even, and (4.2) (resp., (4.6)) if r is odd.*

We have seen in the proof of Theorem 4.1 that the set Pal_r can be identified with the set $\mathcal{M}_{r/2}$ in (3.2), if r is even, or $\widetilde{\mathcal{M}}_{(r-1)/2}$ in (3.3) if r is odd. The only restriction for r here is that $r < n$, which is equivalent to $s < n/2$. Then Theorem 4.1 allows us to conclude that, provided that $s < n/2$, the bounds obtained in Proposition 3.2 (a)–(b) are the dimensions of the sets \mathcal{M}_s and $\widetilde{\mathcal{M}}_s$.

COROLLARY 1. *If \mathcal{M}_s and $\widetilde{\mathcal{M}}_s$ are as in (3.2) and (3.3), respectively, and $s < n/2$, then*

- (a) $\dim \mathcal{M}_s = s(3n - 2s)$, and
- (b) $\dim \widetilde{\mathcal{M}}_s = s(3n - 2s - 1) + n$.

4.2. \top -alternating pencils. Here we provide the counterpart of Theorems 4.1 and 4.2 for \top -alternating pencils.

THEOREM 4.4. (The set of \top -even pencils with bounded rank). *Let r be an integer with $0 \leq r < n$. The set Even_r is an irreducible algebraic set with dimension*

$$\dim \text{Even}_r = \begin{cases} \frac{r}{2} \cdot (3n - r), & \text{if } r \text{ is even,} \\ \frac{r-1}{2} \cdot (3n - r) + n, & \text{if } r \text{ is odd.} \end{cases}$$

Moreover, if r is even, then Even_r is the closure of the congruence orbit of

$$(4.7) \quad K_E^e(\lambda) := \text{diag}(\overbrace{M_{\alpha+1}^b(\lambda), \dots, M_{\alpha+1}^b(\lambda)}^s, \overbrace{M_\alpha^b(\lambda), \dots, M_\alpha^b(\lambda)}^{n-r-s}),$$

where $\frac{r}{2} = (n-r)\alpha + s$ is the Euclidean division of $r/2$ by $n-r$. If r is odd, then Even_r is the closure of the congruence orbit of

$$(4.8) \quad K_E^o(\lambda) := \text{diag}(1, K_E^e(\lambda)),$$

with $K_E^e(\lambda)$ as in (4.7), but now $\frac{r-1}{2} = (n-r)\alpha + s$ is the Euclidean division of $(r-1)/2$ by $n-r$.

Proof. Let \mathcal{C}_{+1} and \mathcal{C}_{-1} be the Cayley transforms in the set of matrix polynomials defined by

$$(4.9) \quad \mathcal{C}_{-1}(Q)(\lambda) = (1 + \lambda)Q\left(\frac{\lambda - 1}{1 + \lambda}\right) \quad \text{and} \quad \mathcal{C}_{+1}(Q)(\lambda) = (1 - \lambda)Q\left(\frac{1 + \lambda}{1 - \lambda}\right),$$

where $Q(\lambda)$ is any matrix polynomial (see [22]). It is straightforward to see that $\mathcal{C}_{+1}(K_E^e) = K_P^e$, and $\mathcal{C}_{+1}(K_E^o) = K_P^o$, with K_P^e and K_P^o being as in (4.1) and (4.2), respectively. Note that, for a given pencil $A_0 + \lambda A_1$ we have

$$\mathcal{C}_{-1}(A_0 + \lambda A_1) = A_0 - A_1 + \lambda(A_0 + A_1), \quad \mathcal{C}_{+1}(A_0 + \lambda A_1) = A_0 + A_1 + \lambda(A_1 - A_0).$$

In particular, $P(\lambda)$ is \mathbb{T} -palindromic if and only if $\mathcal{C}_{+1}(P)$ is \mathbb{T} -even (see also [22, Th. 2.7]). From the definition of \mathcal{C}_{+1} and \mathcal{C}_{-1} is clear that both maps preserve the rank, that is $\text{rank } \mathcal{C}_{+1}(A_0 + \lambda A_1) = \text{rank } \mathcal{C}_{-1}(A_0 + \lambda A_1) = \text{rank } A_0 + \lambda A_1$, for any matrix pencil $A_0 + \lambda A_1$. Moreover, $\mathcal{C}_{-1}(\mathcal{C}_{+1})(A_0 + \lambda A_1) = 2(A_0 + \lambda A_1)$. Therefore, $\mathcal{C}_{+1} : \text{Pal}_r \rightarrow \text{Even}_r$ is an isomorphism of algebraic sets [29, Def. in p. 29]. As a consequence, $\dim \text{Pal}_r = \dim \text{Even}_r$ [29, Cor. 2, Ch. II, §1.3, p. 88] and Even_r is an irreducible algebraic set.

It is also immediate by definition of \mathcal{C}_{+1} that $\mathcal{C}_{+1}(\mathcal{O}_c(K_E^e)) = \mathcal{O}_c(K_P^e)$ and $\mathcal{C}_{+1}(\mathcal{O}_c(K_E^o)) = \mathcal{O}_c(K_P^o)$. Since \mathcal{C}_{+1} is an isomorphism, $\dim \mathcal{C}_{+1}(\mathcal{O}_c(K_E^e)) = \dim \mathcal{O}_c(K_P^e)$ and $\dim \mathcal{C}_{+1}(\mathcal{O}_c(K_E^o)) = \dim \mathcal{O}_c(K_P^o)$. Therefore, Theorem 4.1 implies that the dimensions of $\mathcal{O}_c(K_E^e)$ and $\mathcal{O}_c(K_E^o)$ are as claimed in the statement.

Since $\mathcal{O}_c(K_E^e) \subseteq \text{Even}_r$ and $\mathcal{O}_c(K_E^o) \subseteq \text{Even}_r$, and both $\mathcal{O}_c(K_E^e)$ and $\mathcal{O}_c(K_E^o)$ have the same dimension as Even_r , depending on whether r is even or odd, the result for the \mathbb{T} -even structure follows. \square

THEOREM 4.5. (The set of \mathbb{T} -odd pencils with bounded rank). *Let r be an integer with $0 \leq r < n$. The set Odd_r is an irreducible algebraic set with dimension*

$$\dim \text{Odd}_r = \begin{cases} \frac{r}{2} \cdot (3n - r), & \text{if } r \text{ is even,} \\ \frac{r-1}{2} \cdot (3n - r) + n, & \text{if } r \text{ is odd.} \end{cases}$$

Moreover, if r is even, then Odd_r is the closure of the congruence orbit of

$$(4.10) \quad K_O^e(\lambda) := \text{diag}(\overbrace{M_{\alpha+1}^{-b}(\lambda), \dots, M_{\alpha+1}^{-b}(\lambda)}^s, \overbrace{M_{\alpha}^{-b}(\lambda), \dots, M_{\alpha}^{-b}(\lambda)}^{n-r-s}),$$

where $\frac{r}{2} = (n-r)\alpha + s$ is the Euclidean division of $r/2$ by $n-r$. If r is odd, then Odd_r is the closure of the congruence orbit of

$$(4.11) \quad K_O^o(\lambda) := \text{diag}(\lambda, K_O^e(\lambda)),$$

with $K_O^e(\lambda)$ as in (4.10), but now $\frac{r-1}{2} = (n-r)\alpha + s$ is the Euclidean division of $(r-1)/2$ by $n-r$.

Proof. The result is an immediate consequence of Theorem 4.4, since a matrix pencil $A_0 + \lambda A_1$ is \mathbb{T} -odd if and only if its reversal $A_1 + \lambda A_0$ is \mathbb{T} -even. \square

As for the previous structures, Theorems 4.4 and 4.5 give the generic canonical form of \mathbb{T} -alternating $n \times n$ pencils with rank at most r .

COROLLARY 4.6. (Generic canonical form of \mathbb{T} -alternating pencils with bounded rank). *Let $0 \leq r < n$. The generic canonical structure of $n \times n$ \mathbb{T} -even (respectively, \mathbb{T} -odd) $n \times n$ matrix pencils with rank at most r is (4.7) (resp., (4.10)) if r is even, and (4.8) (resp., (4.11)) if r is odd.*

4.3. Connection with the full rank and non structured cases. The case $n = r$, where the matrix pencils are allowed to be of full rank, deserves some comment. In this case, the generic canonical form for pencils enjoying any of the structures considered in the paper does not contain singular blocks at all. The generic canonical

form of $n \times n$ \mathbb{T} -palindromic pencils can be found in [6, Th. 6]. For the remaining structures, the canonical form can be obtained by applying either the transformation $\lambda \mapsto -\lambda$ (for the \mathbb{T} -anti-palindromic structure, as in the proof of Theorem 4.2), the Cayley transformations \mathcal{C}_{+1} and \mathcal{C}_{-1} (for the \mathbb{T} -even structure, as in the proof of Theorem 4.4), or these Cayley maps followed by reverting the coefficients A_0 and A_1 (for the \mathbb{T} -odd structure, as in the proof of Theorem 4.5). We note that, though Theorem 6 in [6] is stated for strict equivalence (in terms of the KCF) instead of congruence, the generic canonical form (for congruence) can be obtained from that one by gathering each couple of blocks $(\lambda + \mu_i) \oplus (\lambda + 1/\mu_i)$ in the form $\begin{bmatrix} 0 & \lambda + \mu_i \\ 1 + \lambda\mu_i & 0 \end{bmatrix}$.

As a consequence, the generic canonical form in the full-rank case has nothing to do with the generic canonical forms obtained in (4.1)–(4.2), (4.5)–(4.8), and (4.10)–(4.11) for rank-deficient cases. Despite this fact, the formulas for the dimension of Pal_r , Apal_r , Even_r , and Odd_r in Theorems 4.1, 4.2, 4.4, and 4.5, are still valid for $r = n$. To see this, note that by replacing $r = n$ in these formulas we end up, in all cases, with n^2 , which is the dimension of the set of structured $n \times n$ pencils, for any of the structures considered.

It is also interesting to compare the results on the dimension of sets of low-rank structured pencils with the case of non-structured pencils. If we denote by \mathbb{P}_r the set of (unstructured) $n \times n$ matrix pencils with rank at most r , then, as mentioned before, we have

$$\dim \text{Pal}_n = \dim \text{Apal}_n = \dim \text{Even}_n = \dim \text{Odd}_n = \frac{1}{2} \dim \mathbb{P}_n = n^2.$$

One may wonder whether these identities are still true or not for $r < n$. The answer is given in the following corollary.

COROLLARY 2. *The following identities hold, for $r \leq n$:*

$$\dim \text{Pal}_r = \dim \text{Apal}_r = \dim \text{Even}_r = \dim \text{Odd}_r = \begin{cases} \frac{1}{2} \dim \mathbb{P}_r, & \text{if } r \text{ is even,} \\ \frac{1}{2} \dim \mathbb{P}_r - \frac{1}{2}(n - r), & \text{if } r \text{ is odd.} \end{cases}$$

Proof. The result is a direct consequence of the first claim in Theorems 4.1, 4.2, 4.4, and 4.5, together with the fact that $\dim \mathbb{P}_r = r(3n - r)$ [5, Th. 3.3]. \square

The identity for the case r odd in Corollary 2 may be surprising when compared with the case r even. However, the differences between these two cases can be explained by looking at the generic forms provided in (4.1)–(4.2), (4.5)–(4.6) (for the palindromic structures) and (4.7)–(4.8), (4.10)–(4.11) (for the alternating structures). More precisely, in the case r odd, the generic canonical form contains some regular part in all structures, whereas in the case r even it consists entirely of singular blocks. The presence of such regular part imposes some additional restrictions which should lead one to expect some differences in the dimension count. Nonetheless, a full explanation of these particular differences would require to analyze more in detail the algebraic restrictions imposed by the presence of these blocks, something which is beyond the scope of this paper.

5. Conclusions and future work. We have proved that the algebraic sets of \mathbb{T} -palindromic, \mathbb{T} -anti-palindromic, \mathbb{T} -even, and \mathbb{T} -odd matrix pencils with rank at most $r < n$ are irreducible algebraic sets. This is in stark contrast with the case of $n \times n$ unstructured matrix pencils with rank at most r , which is an algebraic set with $r + 1$ irreducible components. We have described these sets of structured matrix pencils with bounded rank as the closures of the congruence orbit of certain structured

pencil given in canonical form. As a consequence, we have determined the generic canonical form of structured pencils with rank at most r , for any of the previous structures. We have also computed the dimension of each of these sets.

A natural continuation of this work is to address the same questions for other structures arising usually in applications, like the Hermitian, skew-Hermitian, $*$ -palindromic, $*$ -anti-palindromic, or $*$ -alternating structures. The sets of $n \times n$ structured pencils satisfying any of these structures are not algebraic sets over \mathbb{C} , but over \mathbb{R} , and for this reason we have not considered them here. Moreover, the description of low-rank pencils with these structures provided in [10] as a sum of rank-1 pencils suggests that the treatment of these structures deserves some additional effort.

Another possible line of research is to extend the results in the paper to matrix polynomials of higher degree. There are some recent contributions in this direction. In [11] the authors have described the generic scalar spectral information of arbitrary (non-structured) matrix polynomials with bounded rank and fixed degree, and in [12] they have obtained an analogous description for the set of skew-symmetric matrix polynomials with bounded rank and fixed degree.

Acknowledgements. This work was done while the author was visiting the TU Berlin within the research group on Numerical Mathematics. The hospitality of the whole group is gratefully acknowledged. The author wishes to acknowledge fruitful and stimulating discussions with C. Mehl and V. Mehrmann. He also thanks J. M. Landsberg for insightful ideas that allowed to prove Proposition 3.2, J. Brundan for pointing out reference [21], and two anonymous referees for the careful reading of the original manuscript, and for very useful comments that led to a significant improvement.

REFERENCES

- [1] L. BATZKE, *Generic rank-one perturbations of structured regular matrix pencils*, Linear Algebra Appl., 458 (2014), pp. 638–670.
- [2] L. BATZKE, *Generic Low-Rank Perturbations of Structured Regular Matrix Pencils and Structured Matrices*, PhD thesis, TU Berlin, Berlin, 2015.
- [3] F. DE TERÁN, *Canonical forms for congruence of matrices: a tribute to H. W. Turnbull and A. C. Aitken*, SeMA Journal, 73 (2016), pp. 7–16.
- [4] F. DE TERÁN AND F. M. DOPICO, *Low rank perturbation of Kronecker structures without full rank*, SIAM J. Matrix Anal. Appl., 29 (2007), pp. 496–529.
- [5] F. DE TERÁN AND F. M. DOPICO, *A note on generic Kronecker orbits of matrix pencils with fixed rank*, SIAM J. Matrix Anal. Appl., 30 (2008), pp. 491–498.
- [6] F. DE TERÁN AND F. M. DOPICO, *The solution of the equation $XA + AX^T = 0$ and its application to the theory of orbits*, Linear Algebra Appl., 434 (2011), pp. 44–67.
- [7] F. DE TERÁN AND F. M. DOPICO, *Generic change of the partial multiplicities of regular matrix pencils under low rank perturbations*, SIAM J. Matrix Anal. Appl., 37 (2016), pp. 823–835.
- [8] F. DE TERÁN, F. M. DOPICO, AND J. M. LANDSBERG, *An explicit description of the irreducible components of the set of matrix pencils with bounded normal rank*, Linear Algebra Appl., 520 (2017), pp. 80–103.
- [9] F. DE TERÁN, F. M. DOPICO, AND J. MORO, *Low rank perturbation of Weierstrass structure*, SIAM J. Matrix Anal. Appl., 30 (2008), pp. 538–547.
- [10] F. DE TERÁN, C. MEHL, AND V. MEHRMANN, *Low rank perturbation of regular structured matrix pencils*, In preparation, (2017).
- [11] A. DMYTRYSHYN AND F. M. DOPICO, *Generic complete eigenstructures for sets of matrix polynomials with bounded rank and degree*, Linear Algebra Appl., 535 (2017), pp. 213–230.
- [12] A. DMYTRYSHYN AND F. M. DOPICO, *Generic skew-symmetric matrix polynomials with fixed rank and fixed odd grade*, Linear Algebra Appl., 536 (2018), pp. 1–18.
- [13] A. DMYTRYSHYN AND B. KÅGSTRÖM, *Orbit closure hierarchies for skew-symmetric matrix pencils*, SIAM J. Matrix Anal. Appl., 35 (2014), pp. 1429–1443.

- [14] A. DMYTRYSHYN, B. KÅGSTRÖM, AND V. V. SERGEICHUK, *Skew-symmetric matrix pencils: Codimension counts and the solution of a pair of matrix equations*, Linear Algebra Appl., 438 (2013), pp. 3375–3396.
- [15] A. DMYTRYSHYN, B. KÅGSTRÖM, AND V. V. SERGEICHUK, *Symmetric matrix pencils: Codimension counts and the solution of a pair of matrix equations*, Electron J. Linear Algebra, 27 (2014), pp. 1–18.
- [16] A. EDELMAN, E. ELMROTH, AND B. KÅGSTRÖM, *A geometric approach to perturbation theory of matrices and matrix pencils. Part I: Versal deformations*, SIAM J. Matrix Anal. Appl., 18 (1997), pp. 653–692.
- [17] A. EDELMAN, E. ELMROTH, AND B. KÅGSTRÖM, *A geometric approach to perturbation theory of matrices and matrix pencils. Part II: A stratification-enhanced staircase algorithm*, SIAM J. Matrix Anal. Appl., 20 (1999), pp. 667–699.
- [18] E. EICH-SOELLNER AND C. FUHRER, *Numerical Methods in Multibody Dynamics*, European Consortium for Mathematics in Industry, Teubner Verlag, 1998.
- [19] F. R. GANTMACHER, *The Theory of Matrices*, Chelsea Publishing Co., 1959.
- [20] R. A. HORN AND C. R. JOHNSON, *Topics in Matrix Analysis*, Cambridge University Press, 1994.
- [21] J. E. HUMPHREYS, *Linear Algebraic Groups*, Springer Verlag, 1975.
- [22] D. S. MACKEY, N. MACKEY, C. MEHL, AND V. MEHRMANN, *Structured polynomial eigenvalue problems: good vibrations from good linearizations*, SIAM J. Matrix Anal. Appl., 28 (2006), pp. 1029–1051.
- [23] D. S. MACKEY, N. MACKEY, C. MEHL, AND V. MEHRMANN, *Jordan structures of alternating matrix polynomials*, Linear Algebra Appl., 432 (2010), pp. 867–891.
- [24] C. MEHL, V. MEHRMANN, AND M. WOJTYLAK, *On the distance to singularity via low rank perturbations*, Oper. Matrices, 9 (2015), pp. 733–772.
- [25] C. MEHL, V. MEHRMANN, AND M. WOJTYLAK, *Parameter-dependent rank-one perturbations of singular Hermitian or symmetric pencils*, SIAM J. Matrix Anal. Appl., 38 (2017), pp. 72–95.
- [26] D. MUMFORD, *Algebraic Geometry (I)*, Springer Verlag, 1995.
- [27] R. RIAZA, *Differential-algebraic systems. Analytical aspects and circuit applications*, World Scientific Publishing Co., 2008.
- [28] C. SCHRÖDER, *A structured Kronecker form for the palindromic eigenvalue problem*, in PAMM, Proc. Appl. Math. Mech. 6, GAMM, ed., Wiley, 2006, pp. 721–722.
- [29] I. R. SHAFAREVICH, *Basic Algebraic Geometry (I)*, Springer Verlag, 2nd Ed., 1994.